

DPP

DAILY PRACTICE PROBLEMS

CLASS : XIth
DATE :

SUBJECT : MATHS
DPP NO. :1

Topic :-RELATIONS AND FUNCTIONS

- Let $A = \{x : -1 \leq x \leq 1\}$ and $f : A \rightarrow A$ such that $f(x) = x|x|$, then f is
 - A bijection
 - Injective but not surjective
 - Surjective but not injective
 - Neither injective nor surjective
- The domain of the function $\sin^{-1}\left(\log_2 \frac{x^2}{2}\right)$ is
 - $[-1, 2] - \{0\}$
 - $[-2, 2] - \{-1, 1\}$
 - $[-2, 2] - \{0\}$
 - $[1, 2]$
- If $f(x) = ax + b$ and $g(x) = cx + d$, then $f\{g(x)\} = g\{f(x)\}$ is equivalent to
 - $f(a) = f(c)$
 - $f(b) = g(b)$
 - $f(d) = g(b)$
 - $f(c) = g(a)$
- The period of the function $f(x) = \sin^4 3x + \cos^4 3x$ is
 - $\pi/2$
 - $\pi/3$
 - $\pi/6$
 - None of these
- Given $f(x) = \log_{10} \left(\frac{1+x}{1-x}\right)$ and $g(x) = \frac{3x+x^3}{1+3x^2}$, then $f \circ g(x)$ equals
 - $-f(x)$
 - $3f(x)$
 - $[f(x)]^3$
 - None of these
- Which of the following functions is not an injective map(s)?
 - $f(x) = |x + 1|, x \in [-1, \infty)$
 - $g(x) = x + \frac{1}{x}, x \in (0, \infty)$
 - $h(x) = x^2 + 4x - 5, x \in (0, \infty)$
 - $h(x) = e^{-x}, x \in [0, \infty)$
- If $f: R \rightarrow R$ and $g: R \rightarrow R$ are defined by $f(x) = x - [x]$ and $g(x) = [x]$ for $x \in R$, where $[x]$ is the greatest integer not exceeding x , then for every $x \in R$, $f(g(x))$ is equal to
 - x
 - 0
 - $f(x)$
 - $g(x)$
- The domain of definition of $f(x) = \sqrt{\frac{\log_{0.3}|x-2|}{|x|}}$, is
 - $[1, 2) \cup (2, 3]$
 - $[1, 3]$
 - $R - (1, 3]$
 - None of these
- $f: R \rightarrow R$ given by $f(x) = 5 - 3 \sin x$, is
 - One-one
 - Onto
 - One-one and onto
 - None of these
- If $f(x + 2y, x - 2y) = xy$, then $f(x, y)$ equals
 - $\frac{x^2 - y^2}{8}$
 - $\frac{x^2 - y^2}{4}$
 - $\frac{x^2 + y^2}{4}$
 - $\frac{x^2 - y^2}{2}$

11. If $f: R \rightarrow R$ is defined as $f(x) = (1-x)^{1/3}$, then $f^{-1}(x)$ is
 a) $(1-x)^{-1/3}$ b) $(1-x)^3$ c) $1-x^3$ d) $1-x^{1/3}$
12. If $f(x+2y, x, x-2y) = xy$, then $f(x, y)$ equals
 a) $\frac{x^2-y^2}{8}$ b) $\frac{x^2-y^2}{4}$ c) $\frac{x^2+y^2}{4}$ d) $\frac{x^2-y^2}{2}$
13. Let $f: [4, \infty[\rightarrow [4, \infty[$ be defined by $f(x) = 5^{x(x-4)}$ then $f^{-1}(x)$
 a) $2 - \sqrt{4 + \log_5 x}$ b) $2 + \sqrt{4 + \log_5 x}$ c) $\left(\frac{1}{5}\right)^{x(x-4)}$ d) Not defined
14. If $f: [2, 3] \rightarrow R$ is defined by $f(x) = x^3 + 3x - 2$, then the range $f(x)$ is contained in the interval
 a) $[1, 12]$ b) $[12, 34]$ c) $[35, 50]$ d) $[-12, 12]$
15. The period of $\sin^2 \theta$, is
 a) π^2 b) π c) 2π d) $\pi/2$
16. If $n \in N$, and the period of $\frac{\cos nx}{\sin(\frac{x}{n})}$ is 4π , then n is equal to
 a) 4 b) 3 c) 2 d) 1
17. For real x , let $f(x) = x^3 + 5x + 1$, then
 a) f is one-one but not onto R b) f is onto R but not one-one
 c) f is one-one and onto R d) f is neither one-one nor onto R
18. The range of the function $f(x) = \frac{1}{2 - \cos 3x}$, is
 a) $[-1/3, 0]$ b) R c) $[1/3, 1]$ d) None of these
19. Let $A = \{2, 3, 4, 5, \dots, 16, 17, 18\}$. Let be the equivalence relation on $A \times A$, cartesian product of A and A , defined by $(a, b) \approx (c, d)$ if $ad = bc$, then the number of ordered pairs of the equivalence class of $(3, 2)$ is
 a) 4 b) 5 c) 6 d) 7
20. Let n be the natural number. Then, the range of the function $f(n) = 8 - n_{P_{n-4}}$, $4 \leq n \leq 6$, is
 a) $\{1, 2, 3, 4\}$ b) $\{1, 2, 3, 4, 5, 6\}$ c) $\{1, 2, 3\}$ d) $\{1, 2, 3, 4, 5\}$