

DPP

DAILY PRACTICE PROBLEMS

CLASS : XIth

DATE :

Solutio

SUBJECT : CHEMISTRY

DPP No. : 1

Topic :- STRUCTURE OF ATOM

- 1 (c)
Isoelectronic species have same number of electron. Mg^{2+} and Na^+ both have 10 electrons hence, they are isoelectronic species.
- 2 (c)
This is obtained by the solution of Schrodinger wave equation
Probability = $\Psi^2 dV$
1st orbital is spherically symmetrical
 $\therefore V = \frac{4}{3}\pi r^3, \therefore \frac{dV}{dr} = 4\pi r^2$
 \therefore Probability = $\Psi^2 4\pi r^2 dr$
- 4 (a)
 $\frac{\Delta E}{(\text{eV})} = \frac{12375}{\lambda_{\text{in } \text{\AA}}} = \frac{12375}{5890} = 2.10 \text{ eV}$
- 5 (b)
1 eV = 1.602×10^{-12} erg.
- 6 (b)
s can have only two values +1/2 and -1/2.
- 7 (c)
The de-Broglie wavelength associated with the charged particle as
For electron, $\lambda = \frac{12.27}{\sqrt{V}} \text{\AA}$
For proton, $\lambda = \frac{0.286}{\sqrt{V}} \text{\AA}$
For α -particles, $\lambda = \frac{0.101}{\sqrt{V}} \text{\AA}$
- 8 (b)
 $\lambda = \frac{h}{mv} = \frac{6.63 \times 10^{-34}}{1.67 \times 10^{-27} \times 1 \times 10^3} = 3.97 \times 10^{-10} \text{ m} \sim 0.40 \text{ nm.}$
- 9 (b)
The number of waves in an orbit = n .
- 10 (a)
 $E \propto \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$
or $E \propto \frac{1}{n^2}$
- 11 (b)
 n is an integer except zero.
- 12 (c)
According to aufbau principle, electrons enter into orbitals according to their energy. The electrons first enters into orbital having lesser value of $(n + l)$. If the value of $n + l$ is same for

two orbitals then the electron will first enter into orbital having lesser value of n .

$$n = 5, l = 0 \therefore n + l = 5 + 0 = 5$$

For other,

$$n = 3, l = 2 \therefore n + l = 3 + 2 = 5$$

\therefore Both of the orbitals have same value for $n + l$.

\therefore Electron will enter into orbital having lower value of n .

\therefore Electron will enter into $n = 3, l = 2$ orbital.

13

(b)

$E = \frac{hc}{\lambda}$, h and c for both causes are same so,

$$\frac{E_1}{E_2} = \frac{\lambda_2}{\lambda_1} = \frac{16000}{8000}$$

$$E_1 = 2E_2$$

14

(c)

When $n = 3$, number of values of l are 0 to $(n - 1)$ i. e., 0, 1, 2

Hence,

when $n = 3$, then $l = 3$ does not exist.

15

(c)

We know that,

$$\Delta E = hc.R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$$

For lowest energy, of the spectral line in Lyman series, $n_1 = 1, n_2 = 2$

Hence,

$$\Delta E = hc.R \left[\frac{1}{1^2} - \frac{1}{2^2} \right]$$

$$\Delta E = \frac{3hcR}{4}$$

16

(c)

Cathode rays are fastly moving electrons.

17

(c)

$$1. \quad n = 4, l = 0, m = 0, s = +\frac{1}{2}$$

$\rightarrow 4s$ energy level.

$$2. \quad n = 3, l = 1, m = -1, s = +\frac{1}{2}$$

$\rightarrow 3p$ energy level.

$$3. \quad n = 3, l = 2, m = -2, s = +\frac{1}{2}$$

$\rightarrow 3d$ energy level.

$$4. \quad n = 3, l = 0, m = 0, s = +\frac{1}{2}$$

$\rightarrow 3s$ energy level.

According to aufbau principle, the energy of orbitals (other than H-atom) depend upon $n + 1$ value.

$$n + l \text{ for } 3d = 3 + 2 = 5$$

So, it is highest energy level (in the given options).

18

(d)

Each one possesses mass.

19

(c)

X-rays have larger wavelength than γ -rays.

20

(c)

$$\Delta E = \frac{hc}{\lambda}$$

ANSWER-KEY										
Q.	1	2	3	4	5	6	7	8	9	10
A.	C	D	B	C	C	B	B	C	B	C
Q.	11	12	13	14	15	16	17	18	19	20
A.	D	B	B	D	A	C	B	B	C	A

**SMARTLEARN
COACHING**