

DPP

DAILY PRACTICE PROBLEMS

CLASS : XIth
DATE :

SUBJECT : MATHS
DPP NO. :2

Topic :-RELATIONS AND FUNCTIONS

1. Let X and Y be subsets of R , the set of all real numbers. The function $f: X \rightarrow Y$ defined by $f(x) = x^2$ for $x \in X$ is one-one but not onto, if (Here, R^+ is the set of all positive real numbers)
 - a) $X = Y = R^+$
 - b) $X = R, Y = R^+$
 - c) $X = R^+, Y = R$
 - d) $X = Y = R$
2. If $f(x) \cdot f(1/x) = f(x) + f(1/x)$ and $f(4) = 65$, then $f(6)$ is
 - a) 65
 - b) 217
 - c) 215
 - d) 64
3. The graph of the function of $y = f(x)$ is symmetrical about the line $x = 2$, then
 - a) $f(x+2) = f(x-2)$
 - b) $f(2+x) = f(2-x)$
 - c) $f(x) = f(-x)$
 - d) $f(x) = -f(-x)$
4. If $f(x) = \begin{cases} -1; & x < 0 \\ 0; & x = 0 \\ 1; & x > 0 \end{cases}$ and $g(x) = x(1-x^2)$, then
 - a) $f \circ g(x) = \begin{cases} -1; & -1 < x < 0 \text{ or } x > 1 \\ 0; & x = 0, 1, -1 \\ 1; & 0 < x < 1 \end{cases}$
 - b) $f \circ g(x) = \begin{cases} -1; & -1 < x < 0 \\ 0; & x = 0, 1, -1 \\ 1; & 0 < x < 1 \end{cases}$
 - c) $f \circ g(x) = \begin{cases} -1; & -1 < x < 0 \text{ or } x > 1 \\ 0; & x = 0, 1, -1 \\ 1; & 0 < x < 1 \text{ or } x < -1 \end{cases}$
 - d) $f \circ g(x) = \begin{cases} 1; & -1 < x < 0 \text{ or } x > 1 \\ 0; & x = 0, 1, -1 \\ 1; & 0 < x < 1 \text{ or } x < -1 \end{cases}$
5. $x_2 = xy$ is a relation which is
 - a) Symmetric
 - b) Reflexive and transitive
 - c) Transitive
 - d) None of these
6. The period of $f(x) = \sin\left(\frac{\pi x}{n-1}\right) + \cos\left(\frac{\pi x}{n}\right)$, $n \in Z$, $n > 2$, is
 - a) $2n \pi(n-1)$
 - b) $4(n-1)\pi$
 - c) $2n(n-1)$
 - d) None of these
7. $f: [-4,0] \rightarrow R$ is given by $f(x) = e^x + \sin x$, its even extension to $[-4,4]$, is
 - a) $-e^{-|x|} - \sin|x|$
 - b) $e^{-|x|} - \sin|x|$
 - c) $e^{-|x|} + \sin|x|$
 - d) $-e^{-|x|+\sin|x|}$
8. Let $f: R \rightarrow R$ be a function defined by $f(x) = -\frac{|x|^3+|x|}{1+x^2}$, then the graph of $f(x)$ lies in the
 - a) I and II quadrants
 - b) I and III quadrants
 - c) II and III quadrants
 - d) III and IV quadrants

9. The domain of the real valued function $f(x) = \sqrt{1 - 2x} + 2 \sin^{-1}\left(\frac{3x-1}{2}\right)$ is
 a) $\left[-\frac{1}{3}, 1\right]$ b) $\left[\frac{1}{2}, 1\right]$ c) $\left[-\frac{1}{2}, \frac{1}{3}\right]$ d) $\left[-\frac{1}{3}, \frac{1}{2}\right]$
10. The domain of function $f(x) = \log_{(x+3)}(x^2 - 1)$ is
 a) $(-3, -1) \cup (1, \infty)$
 b) $[-3, -1] \cup [1, \infty)$
 c) $(-3, -2) \cup (-2, -1) \cup (1, \infty)$
 d) $[-3, -2] \cup (-2, -1) \cup [1, \infty)$
11. The range of the function $f(x) = x^2 - 6x + 7$ is
 a) $(-\infty, 0)$ b) $[-2, \infty)$ c) $(-\infty, \infty)$ d) $(-\infty, -2)$
12. The inverse of the function $f: R \rightarrow (-1, 3)$ is given by $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}} + 2$
 a) $\log\left(\frac{x-1}{x+1}\right)^{-2}$ b) $\log\left(\frac{x-2}{x-1}\right)^{1/2}$ c) $\log\left(\frac{x}{2-x}\right)^{1/2}$ d) $\log\left(\frac{x-1}{3-x}\right)^{1/2}$
13. If $f(x) = \frac{4^x}{4^x + 2}$, then $f\left(\frac{1}{97}\right) + f\left(\frac{2}{97}\right) + \dots + f\left(\frac{96}{97}\right)$ is equal to
 a) 1 b) 48 c) -48 d) -1
14. The period of the function $f(x) = \frac{\sin 8x \cos x - \sin 6x \cos 3x}{\cos 2x \cos x - \sin 3x \sin 4x}$ is
 a) π b) 2π c) $\frac{\pi}{2}$ d) None of these
15. Let $f: R \rightarrow R: f(x) = x^2$ and $g: R \rightarrow R: g(x) = x + 5$, then gof is
 a) $(x + 5)$ b) $(x + 5^2)$ c) $(x^2 + 5^2)$ d) $(x^2 + 5)$
16. The function $f(x) = \log_{2x-5}(x^2 - 3x - 10)$ is defined for all x belonging to
 a) $[5, \infty)$ b) $(5, \infty)$ c) $(-\infty, +5)$ d) None of these
17. Range of the function $f(x) = \frac{x^2}{x^2 + 1}$ is
 a) $(-1, 0)$ b) $(-1, 1)$ c) $[0, 1)$ d) $(1, 1)$
18. Let $f(x) = |x - 1|$. Then,
 a) $f(x^2) = [f(x)]^2$
 b) $f(|x|) = |f(x)|$
 c) $f(x + y) = f(x) + f(y)$
 d) None of these
19. If $f(x) = a^x$, which of the following equalities do not hold?
 a) $f(x + 2) - 2f(x + 1) + f(x) = (a - 1)^2 f(x)$
 b) $f(-x)f(x) - 1 = 0$
 c) $f(x + y) = f(x)f(y)$
 d) $f(x + 3) - 2f(x + 2) + f(x + 1) = (a - 2)^2 f(x + 1)$
20. Let $A = \{x \in R: x \leq 1\}$ and $f: A \rightarrow A$ be defined as $f(x) = x(2 - x)$. Then, $f^{-1}(x)$ is

a) $1 + \sqrt{1 - x}$

b) $1 - \sqrt{1 - x}$

c) $\sqrt{1 - x}$

d) $1 \pm \sqrt{1 - x}$



**SMARTLEARN
COACHING**