

## DPP

DAILY PRACTICE PROBLEMS

CLASS : XI<sup>th</sup>  
DATE :

SUBJECT : MATHS  
DPP NO. : 2

### Topic :- COMPLEX NUMBERS AND QUADRATIC EQUATIONS

- The number of integral solutions of  $2(x + 2) > x^2 + 1$ , is  
 a) 2                                      b) 3                                      c) 4                                      d) 5
- If one root of the equation  $(a - b)x^2 + ax + 1 = 0$  be double the other and if  $a \in R$ , then the greatest value of  $b$  is  
 a)  $9/8$                                       b)  $7/8$                                       c)  $8/9$                                       d)  $8/7$
- The argument of  $(1 - i\sqrt{3})(1 + i\sqrt{3})$  is  
 a)  $60^\circ$                                       b)  $120^\circ$                                       c)  $210^\circ$                                       d)  $240^\circ$
- If the area of the triangle on the complex plane formed by the points  $z, z + iz$ , and  $iz$  is 200, then the value of  $|3z|$  must be equal to  
 a) 20                                      b) 40                                      c) 60                                      d) 80
- If the roots of the equation  $bx^2 + cx + a = 0$  be imaginary, then for all real values of  $x$ , the expression  $3b^2x^2 + 6bcx + 2c^2$  is  
 a) Greater than  $4ab$                       b) Less than  $4ab$                       c) Greater than  $-4ab$                       d) Less than  $-4ab$
- If  $(ax^2 + c)y + (d x^2 + c') = 0$  and  $x$  is a rational function of  $y$  and  $ac$  is negative, then  
 a)  $ac' + a'c = 0$                       b)  $\frac{a}{a'} = \frac{c}{c'}$                       c)  $a^2 + c^2 = a'^2 + c'^2$                       d)  $aa' + cc' = 1$
- If  $n$  is a positive integer, then  $(1 + i\sqrt{3})^n + (1 - i\sqrt{3})^n$  is equal to  
 a)  $2^{n-1} \cos \frac{n\pi}{3}$                       b)  $2^n \cos \frac{n\pi}{3}$                       c)  $2^{n+1} \cos \frac{n\pi}{3}$                       d) None of these
- The points represented by the complex numbers  $1 + i, -2 + 3i, \frac{5}{3}i$  on the argand diagram are  
 a) Vertices of an equilateral triangle                      b) Vertices of an isosceles triangle  
 c) Collinear                                      d) None of the above
- If the amplitude of  $z - 2 - 3i$  is  $\frac{\pi}{4}$ , then the locus of  $z = x + iy$ , is  
 a)  $x + y - 1 = 0$                       b)  $x - y - 1 = 0$                       c)  $x + y + 1 = 0$                       d)  $x - y + 1 = 0$
- The value of  $\frac{[(\cos 20^\circ + i \sin 20^\circ)(\cos 75^\circ + i \sin 75^\circ)(\cos 10^\circ + i \sin 10^\circ)]}{\sin 15^\circ - i \cos 15^\circ}$  is  
 a) 0                                      b) -1                                      c)  $i$                                       d) 1

11. Let  $\alpha, \beta$  be the roots of  $x^2 + bx + 1 = 0$ . Then the equation whose roots are  $-\left(\alpha + \frac{1}{\beta}\right)$  and  $-\left(\beta + \frac{1}{\alpha}\right)$ , is
- a)  $x^2 = 0$                       b)  $x^2 + 2bx + 4 = 0$       c)  $x^2 - 2bx + 4 = 0$       d)  $x^2 - bx + 1 = 0$
12. The vector  $z = -4 + 5i$  is turned counterclockwise through an angle of  $180^\circ$  and stretched  $1\frac{1}{2}$  times. The complex number corresponding to newly obtained vector is
- a)  $6 - \frac{15}{2}i$                       b)  $-6 + \frac{15}{2}i$                       c)  $6 + \frac{15}{2}i$                       d) None of these
13. If  $(3 - i)z = (3 - i)\bar{z}$ , then the complex number  $z$  is
- a)  $a(3 - i), a \in R$               b)  $\frac{a}{(3+i)}, a \in R$               c)  $a(3 + i), a \in R$               d)  $a(-3 + i), a \in R$
14. For real values of  $x$ , the expression  $\frac{(x-b)(x-c)}{(x-a)}$  will assume all real values provided
- a)  $a \leq c \leq b$                       b)  $b \geq a \geq c$                       c)  $b \leq c \leq a$                       d)  $a \geq b \geq c$
15. If  $(x - 1)^3$  is a factor of  $x^4 + ax^3 + bx^2 + cx - 1$ , then the other factor is
- a)  $x - 3$                       b)  $x + 1$                       c)  $x + 2$                       d)  $x - 1$
16. The centre of a square is at the origin and  $1 + i$  is one of its vertices. The extremities of its diagonals which does not pass through this vertex are
- a)  $1 - i, -1 + i$                       b)  $1 - i, -1 - i$                       c)  $-1 + i, -1 - i$                       d) None of these
17. If  $p(x) = ax^2 + bx + c$  and  $Q(x) = -ax^2 + dx + c$ , where  $ac \neq 0$ , then  $P(x)Q(x) = 0$  has at least
- a) Four real roots                      b) Two real roots  
c) Four imaginary roots                      d) None of these
18. If  $a = \cos \theta + i \sin \theta$ , then  $\frac{1+a}{1-a}$  is equal to
- a)  $\cot \frac{\theta}{2}$                       b)  $\cot \theta$                       c)  $i \cot \frac{\theta}{2}$                       d)  $i \tan \frac{\theta}{2}$
19. If  $x^2 + 2ax + b \geq c, \forall x \in R$ , then
- a)  $a - c \geq a^2$                       b)  $c - a \geq b^2$                       c)  $a - b \geq c^2$                       d) None of these
20. Let  $A, B, C$  be three collinear points which are such that  $AB \cdot AC = 1$  and the points are represented in the Argand plane by the complex numbers  $0, z_1$  and  $z_2$  respectively, Then,
- a)  $z_1 z_2 = 1$                       b)  $z_1 \bar{z}_2 = 1$                       c)  $|z_1| |z_2| = 1$                       d) None of these