

CLASS: XIth DATE:

Solutio

SUBJECT: CHEMISTRY

DPP No. : 3

Topic:- SOME BASIC CONCEPTS OF CHEMISTRY

(b) 1

See mole ratio A : B : C :: 1 :: 2 :: 1

2

1 mg C₄H₁₀ =
$$\frac{14N}{58} \times 10^{-3}$$
 atoms,

1 mg N₂ =
$$\frac{2N \times 10^{-3}}{28}$$
 atoms,

$$1 \text{ mg Na} = \frac{N \times 10^{-3}}{23} \text{ atoms,}$$

1 mL = 1 g H₂0 =
$$\frac{3N}{18}$$
 atoms,

(: M g of a substance = N molecules = $a \times N$ atoms; where a is number of atoms in one molecule).

3 (c)

An aromatic hydrocarbon (empirical formula C₅H₄)

 $+H_2SO_4 \rightarrow monosulphonic acid$

v = 0.104 g of monosulphonic acid required 10 mL of $\frac{N}{20}$ NaOH for complete neutralisation

$$\therefore \frac{0.104}{n(5 \times 12 + 4 \times 1)} = \frac{1}{20} \times 10 \times 10^{-3}$$
104

$$n = \frac{104}{32} = 3.25 \approx 3$$

The molecular formula of hydrocarbon will be $C_{15}H_{12}$.

In 12 g carbon, mass of C-14 isotope = $12 \times \frac{2}{100} = 0.24g$

: Number of C-14 atoms in 12 g of
$$C = \frac{0.24}{14} \times 6.02 \times 10^{23}$$

$$= 1.032 \times 10^{22}$$

5 (b)

To prepare 20 g zinc sulphate crystals, zinc required

$$= \frac{22.65}{100} \times 20$$
$$= 4.53g$$

6 (b)

Number of gram molecules
$$=\frac{6.02 \times 10^{25}}{6.02 \times 10^{23}} = 100$$

7 (a)

Ferrous is Fe²⁺

8

$$M = \frac{5}{34 \times 100/1000} = 1.47$$

9 (b)

 4.6×10^{22} atoms weight = 13.8 g

Hence, 6.02×10^{23} atoms will weigh

$$= \frac{13.8 \times 6.02 \times 10^{23}}{4.6 \times 10^{22}} = 108.6 \text{ g (molar mass)}$$

10 (c)

Eq. of HCl = Eq. of $CaCO_3$

Thus,
$$\frac{w}{36.5} = \frac{100}{50}$$
;

50 g HCl is present in 100 g HCl solution and thus, volume of solution required for,

73 g HCl =
$$\frac{73 \times 100}{50}$$
 = 146g.

12 (d)

> The law of constant composition—According to this law, "A chemical compound is always found to be made up of the same elements combined together in the same proportions by weights". This law is same as law of definite proportions.

13 (d)

Atomic weight of the element

$$X = 6.643 \times 10^{-23} \times N_A = 40$$

$$X = 6.643 \times 10^{-23} \times N_A = 40$$

No. of moles of $X = \frac{20 \times 1000}{40} = 500$

14 (a)

Limiting reagent is one which is completely consumed in reaction.

15

ppm = wt. of solute in 10^6 g H_2 0

10³ g H₂O contains 10 g CaCO₃

$$\therefore 10^6 \text{ g H}_2\text{ O contains} = \frac{10 \times 10^6}{10^3} = 10,000 \text{ ppm CaCO}_3$$

16 (d)

$$BaCl_2 + H_2SO_4 \rightarrow BaSO_4 + 2HCl$$

mm
$$20 \times 0.5$$
 20×1

mm formed

Milli mole of
$$BaSO_4 = 10$$

$$r \qquad Mole \text{ of } BaSO_4 = 10^{-2}$$

17 (d)

Percentage of element *M* in $M_2O_3 = 53$

Let the atomic mass of M = x

Mass of
$$M$$
in $M_2O_3 = 2x$

Total atomic mass of
$$M_2O_3 = 2x + 16 \times 3$$

$$= 2x + 48$$

Percentage of an element

Mass of an element in a compound × 100

Total mass of compound

$$53 = \frac{2x}{2x + 48} \times 100$$

$$53(2x + 48) = 200x$$

$$x = 27$$

18 (a)

H₃BO₃ accepts OH⁻ ions to act as weak monobasic Lewis acid.

$$H_3BO_3 + H_2O \rightarrow B(OH)_4^- + H^+; K_a = 10^{-9}$$

19 (a)

Meq. of KOH added =
$$25 \times 0.4210 = 10.525$$

Meq. of KOH left = $8.46 \times 0.2732 \times 2 = 4.623$
∴ Meq. of KOH used by oil = $10.525 - 4.623 = 5.902$
or $\frac{w}{56} \times 1000 = 5.902$
or $\frac{w}{KOH} = 0.3305 \text{ g}$
∴ Saponification no.
= wt. of KOH used in mg per g of oil = $\frac{0.3305}{1.5763} \times 1000$
= $\frac{0.3305}{1.5763} \times 1000$
= $\frac{209.6}{1.5763} \times 1000$
(C) $\frac{(NH_4)_2SO_4}{2SO_4} \rightarrow \frac{2NH_3}{2SO_4} + \frac{2HCl}{2SO_4}$
 $\frac{132}{2SO_4} = \frac{2NH_3}{2SO_4} = \frac{2HCl}{2SO_4}$
 $\frac{132}{2SO_4} = \frac{132\times292}{73} = \frac{2}{2SO_4}$
 $\frac{132\times292}{2SO_4} = \frac{132\times292}{2SO_4} = \frac{1$

ANSWER-KEY										
Q.	1	2	3	4	5	6	7	8	9	10
Α.	В	D	C	A	В	В	A	В	В	С
			A				2			
Q.	11	12	13	14	15	16	17	18	19	20
Α.	A	D	D	A	D	D	D	A	A	C

SMARTLEARN COACHING