

DPP

DAILY PRACTICE PROBLEMS

CLASS : XIth

DATE :

Solutio

SUBJECT : CHEMISTRY

DPP No. : 3

Topic :- STRUCTURE OF ATOM

1

(c)

$$E_n = \frac{E_1}{n^2} \times Z^2$$

$$= \frac{-13.6}{4} \times 9 = -30.6 \text{ eV}$$

(for the excited state, $n = 2$ and for Li^{2+} ion, $Z = 3$)

2

(b)

Given, azimuthal quantum number (l) = 2

Number of orbital's = $(2l+1)$

$$= (2 \times 2 + 1) = 4 + 1 = 5$$

3

(b)

Heaviest atom has mass no. 238, (i. e., ${}_{92}\text{U}^{238}$) and lighter one is ${}_{1}\text{H}^1$.

4

(d)

$$\lambda = \frac{h}{mu}$$

5

(c)

p_x orbital has two lobes on x -axis.

6

(d)

f -orbital has 7 orientations.

8

(b)

III shell is more closer to nucleus.

9

(b)

Ar and Ca^{2+} are isoelectronic species as they have same number of electrons, i. e., 18.

10

(b)

$$p = mu = \frac{h}{\lambda} \text{ and } E = \frac{hc}{\lambda}$$

$$\therefore E = \frac{c}{\lambda} \cdot p \cdot \lambda = c \cdot p$$

11

(a)

$$\Delta x \cdot \Delta v \geq \frac{h}{4\pi m}$$

$$\Delta x \geq \frac{6.62 \times 10^{-34}}{4 \times 3.14 \times 25 \times 10^{-3} \times 10^{-5}}$$

$$= 2.10 \times 10^{-28} \text{ m}$$

12

(d)

Mass of neutron = $1.675 \times 10^{-27} \text{ kg}$.

13

(c)

$$\lambda = \frac{h}{mu} = \frac{6.62 \times 10^{-34}}{66 \times 10^3 \times 1}$$

15

(c)

$n = 4$ (4th shell)

$$l = 2(d\text{-subshell})$$

$$m_1 = -2(d_{xy} \text{ orbital})$$

$$s = +\frac{1}{2}(\uparrow)$$

Hence, electron belongs to 4d-orbital.

16

(d)

The four lobes of $d_{x^2-y^2}$ orbital are lying along x and y axes, while the two lobes of d_{z^2} orbital are lying along z -axis, and contain a ring of negative charge surrounding the nucleus in xy plane
 $2s$ orbitals has one spherical node, where electron density is zero

p -orbital have direction character

Orbital $\rightarrow p_z \quad p_x \quad p_y$

$m \rightarrow 0 \quad \pm 1 \quad \pm 1$

Nodal plane $\rightarrow xy \quad yz \quad zx$

17

(c)

d_{xy} orbital lies at 45° angle in between x -and y -axes.

18

(d)

According to Pauli exclusion principle.

19

(b)

$$E = \frac{hc}{\lambda}$$

20

(d)

Cu has configuration $[\text{Ar}]3d^{10}, 4s^1$; the two electrons are lost, one from $4s^1$ and one from $3d^{10}$.

ANSWER-KEY

Q.	1	2	3	4	5	6	7	8	9	10
A.	C	B	B	D	C	D	D	B	B	B
Q.	11	12	13	14	15	16	17	18	19	20
A.	A	D	C	B	C	D	C	D	B	D

SMARTLEARN
COACHING