

## DPP

DAILY PRACTICE PROBLEMS

CLASS : XI<sup>TH</sup>

DATE :

Solutio

SUBJECT : PHYSICS

DPP NO. : 3

### Topic :- UNITS AND MEASUREMENTS

1

(b)

Given,  $p = \frac{a-t^2}{bx}$  or  $pbx = a - t^2$

By the law of homogeneity of dimensional equation.

Dimensions of  $a$  = dimensions of  $t^2 = [T^2]$

Dimensions of  $b$  = dimensions of  $\frac{t^2}{px} = [M^{-1}T^4]$

So, dimensions of  $\frac{a}{b}$  is  $[MT^{-2}]$ .

2

(d)

$$f = \frac{uv}{u+v}, \frac{\Delta f}{f} = \frac{\Delta u}{u} + \frac{\Delta v}{v} + \frac{(u+v)}{u+v}$$

4

(b)

$$L = \frac{\phi}{I} = \frac{Wb}{A} = \text{Henry}$$

6

(b)

$$r_1 = 10^{-15} \text{m}, r_2 = 10^{26} \text{m}$$

$$\log r = \frac{1}{2} [\log 10^{-15} + \log 10^{26}]$$

$$= \frac{1}{2} [-15 + 26] = 5.5 \approx 6 \Rightarrow r = 10^6 \text{m}$$

7

(d)

The dimensions of  $x$  = dimensions of  $\frac{v_0}{A}$

Therefore, out of the given options  $v_0$  has dimensions equal to  $[M^0LT^{-1}]$  and  $A$  has dimensions equal to  $[M^0L^0T^{-1}]$

$$\text{So, that } \frac{[v_0]}{[A]} = \frac{[M^0LT^{-1}]}{[M^0L^0T^{-1}]} = [L] \\ = \text{dimension of } x$$

8

(c)

$$1 \text{ nm} = 10^{-9} \text{ m} = 10^{-7} \text{ cm}$$

9

(c)

$$\text{Electric potential } V = IR, [R] = \left[ \frac{V}{I} \right] = \left[ \frac{\text{Work done}}{\text{Charge} \times I} \right]$$

$$= \frac{[ML^2T^{-2}]}{[A^2T]} = [ML^2T^{-3}A^{-2}]$$

10

(d)

According to Planck's hypothesis

$$E = hv$$

Or  $h = \frac{E}{v}$

Substituting the dimensions of energy  $E$  and frequency  $v$ , we get

$$[h] = \frac{[ML^2T^{-2}]}{[T^{-1}]}$$



12  $\therefore [h] = [ML^2T^{-1}]$   
**(a)**  
 The dimension of  $y = \frac{e^2}{4\pi\epsilon_0 hc}$   
 Putting the dimensions of  
 $[e] = [Q] = [AT]$   
 $[\epsilon_0] = [M^{-1}L^{-3}T^4A^2], h = [ML^2T^{-1}], c = [LT^{-1}]$   

$$y = \frac{[A^2T^2]}{[M^{-1}L^{-3}T^4A^2][ML^2T^{-1}][LT^{-1}]}$$
  
 $y = [M^0L^0T^0]$

13 **(b)**  
 Volume  $V = l \times b \times t$   
 $= 12 \times 6 \times 2.45 = 176.4 \text{ cm}^3$   
 $V = 1.764 \times 10^2 \text{ cm}^3$   
 Since, the minimum number of significant figure is one in breadth, hence volume will also contain only one significant figure. Hence,  $V = 2 \times 10^2 \text{ cm}^3$

14 **(d)**  
 Percentage error in  
 $A = \left( 2 \frac{\Delta a}{a} + 3 \frac{\Delta b}{b} + \frac{\Delta c}{c} + \frac{1}{2} \frac{\Delta d}{d} \right) \times 100\%$   
 $= 2 \times 1 + 3 \times 3 + 2 + \frac{1}{2} \times 2$   
 $= 2 + 9 + 2 + 1 = 14\%$

16 **(a)**  
 The unit of  $\frac{1}{2} \epsilon E^2 = \frac{C^2}{Nm^2} \left( \frac{N}{C} \right)^2$   
 $= \frac{C^2 N^2}{Nm^2 C^2} = \frac{N}{m^2} = \frac{Nm}{m^3}$   
 $= \frac{J}{m^3} = \text{energy density}$

17 **(d)**  
 $v = at + bt^2$   
 $[v] = [bt^2] \text{ or } LT^{-1} = bT^2 \Rightarrow [b] = [LT^{-3}]$

18 **(b)**  
 $6 \times 10^{-5} = 60 \times 10^{-6} = 60 \text{ microns}$

19 **(b)**  
 Surface tension =  $\frac{\text{Force}}{\text{Length}} = \text{newton/metre}$

20 **(d)**  
 $C = \frac{1}{\sqrt{\mu_0 \epsilon_0}} \Rightarrow \frac{1}{\mu_0 \epsilon_0} = c^2 = [L^2T^{-2}]$

ANSWER-KEY										
Q.	1	2	3	4	5	6	7	8	9	10
A.	B	D	D	B	D	B	D	C	C	D
Q.	11	12	13	14	15	16	17	18	19	20
A.	C	A	B	D	A	A	D	B	B	D



**SMARTLEARN  
COACHING**