

DPP

DAILY PRACTICE PROBLEMS

Class : XIIth
Date :

Solutio


Subject : CHEMISTRY
DPP No. : 1

Topic :- Amines

- 1 (a)
Hofmann bromamide reaction is used to prepare 1° amine from primary amides. In this method, amides are treated with bromine in presence of KOH.

$$R - CONH_2 + 4KOH + Br_2 \xrightarrow{\Delta} RNH_2 + K_2CO_3 + 2KBr + 2H_2O$$
- 4 (d)
The amines are basic in nature due to presence of lone pair of electron on nitrogen. The 2° amines are basic among 1°, 2° and 3° amines because of steric effect and hydration effect
- 5 (a)
This is carbylamine reaction.
- 6 (c)
p-amine forms alcohol; *s*-amine forms only nitrosoamine.
- 7 (a)

$$CH_3NH_2 + C_6H_5SO_2Cl \rightarrow C_6H_5SO_2NHCH_3 \xrightarrow[\text{(alkali)}]{NaOH} C_6H_5SO_2N(Na)CH_3$$

1° amine	hinsberg's	N-methyl benzene	soluble sodium salt
	Reagent	sulphonamide	
- 8 (d)


$p\text{-toluidine} + CHCl_3 + alc\ 3KOH \rightarrow \text{N-cyanotoluidine} + 3KCl + 3H_2O$


it is an example of carbylamine reaction

9 (a)
Cyanides are hydrolysed either by alkali or acid to give carboxylic acid.

$$R - CN + 2H_2O \xrightarrow{NaOH} R - COOH + NH_3$$

10 (a)

$$C_2H_5NH_2 + NOCl \rightarrow C_2H_5Cl + N_2 + H_2O$$


11 (b)
By using H₂S, NH₃ as reagent, selective reduction takes place


12 (b)
2° amines are more basic than 1° amines due to +ve IE of —CH₃ gp. In amide the resonance gives rise to less availability to electron pair for coordination and thus less basic. The negative

charge developed on N in $\text{CH}_3-\text{C} \begin{matrix} \diagup \text{NH} \\ \diagdown \text{NH}_2 \end{matrix}$

due to resonance makes it more basic.

14



Nitration of aniline also gives *m*-nitro aniline in strong acidic medium because in strong acidic condition protonation of $-\text{NH}_2$ group gives anilinium ion ($+\text{NH}_3$) which is deactivating in nature and of *m*-directive nature


15

(a)

Grabriel's synthesis : Phthalimide is reacted with KOH to form potassium phthalimide. The potassium salt is treated with an alkyl halide. The product N-alkyl phthalimide is put to hydrolyse with hydrochloric acid, then primary amine is formed.

16

(b)




Nitrobenzene Aniline

Nitrobenzene in reduction with Sn and HCl produce aniline. Hence, 'X' is identified as $-\text{NH}_2$ group.

17

(d)



Acetaldehyde

ethyl amine

ethyl chloride

(A)

(B)

(C)

19

(b)

$\text{CH}_3\text{CH}_2-\text{O}-\text{N}=\text{O}$ is a nitrite derivative, hence it is not a nitro derivative.

20

(c)

Basic nature of an amine depends upon availability of lone pair on nitrogen atom. If lone pair is easily available the compound would be more basic.

Due to +I effect of methyl group, methyl amine is more basic than ammonia and dimethyl amine is more basic than methyl amine. While aniline is a weaker base than ammonia due to delocalization of lone pair of nitrogen atom at different position.

ANSWER-KEY										
Q.	1	2	3	4	5	6	7	8	9	10
A.	A	A	A	D	A	C	A	D	A	A
Q.	11	12	13	14	15	16	17	18	19	20
A.	B	B	A	C	A	B	D	D	B	C



**SMARTLEARN
COACHING**