

DPP

DAILY PRACTICE PROBLEMS

Class : XIth

Date :

Solutio

Subject : MATHS

DPP No. : 1

Topic :- Applications of Derivatives

1

(b)


We have,

$$f(x) = x(x-1)^2$$

$$\Rightarrow f'(x) = (x-1)^2 + 2x(x-1)$$

$$\Rightarrow f'(x) = (x-1)(3x-1)$$

The changes in the signs of $f'(x)$ are shown in diagram




Clearly, $f(x)$ attains a local maximum at $x = \frac{1}{3}$ and a local minimum at $x = 1$

$$\therefore \text{Maximum value of } f(x) = f\left(\frac{1}{3}\right) = \frac{4}{27}$$

2

(a)

$$\text{Since, } 2\pi k - \frac{\pi}{2} \leq \sin x \leq 2\pi k + \frac{\pi}{2}$$



For $k = 0$

$$-\frac{\pi}{2} < \sin x < \frac{\pi}{2}$$


Which increase from -1 to 1 .

Similarly, for other values of k it is increase from -1 to 1 .

3

(c)

$$\text{Volume of cone, } V = \frac{\pi}{3} r^2 h$$





$$\Rightarrow V = \frac{\pi}{3} r^2 \sqrt{l^2 - r^2}$$

On differentiating w.r.t. r , we get

$$\frac{dV}{dr} = \frac{\pi}{3} \left[2r\sqrt{l^2 - r^2} + \frac{r^2}{2\sqrt{l^2 - r^2}}(-2r) \right]$$

Put $\frac{dV}{dr} = 0$

$$\Rightarrow 2r\sqrt{l^2 - r^2} - \frac{r^3}{\sqrt{l^2 - r^2}} = 0$$

$$\Rightarrow r[2(l^2 - r^2) - r^2] = 0$$

$$\Rightarrow r = \pm l \sqrt{\frac{2}{3}}$$

\therefore At $r = l \sqrt{\frac{2}{3}}$, $\frac{d^2V}{dr^2} < 0$, maxima

$$\therefore h = \sqrt{l^2 - \frac{2}{3}l^2} = \frac{l}{\sqrt{3}}$$

In ΔABC , $\tan\theta = \frac{r}{h} = \frac{l\sqrt{\frac{2}{3}}}{\frac{l}{\sqrt{3}}} = \sqrt{2}$

4

(c)

Given that $f(x) = \sin x - bx + c$

$$\therefore f'(x) = \cos x - b$$

For decreasing, $f'(x) < 0$, for all $x \in R$.

$$\Rightarrow \cos x < b \text{ for all } x \in R \Rightarrow b > 1.$$

5

(c)

Given, $f(x) = 2x^3 + 3x^2 - 12x + 1$

$$\Rightarrow f'(x) = 6x^2 + 6x - 12$$

For $f(x)$ to be decreasing, $f'(x) < 0$

$$\Rightarrow 6(x^2 + x - 2) < 0$$

$$\Rightarrow (x + 2)(x - 1) < 0$$

$$\Rightarrow x \in (-2, 1)$$

6

(a)

We have,

$$f(x) = 2x + \cot^{-1} x + \log(\sqrt{1+x^2} - x)$$

$$\therefore f'(x) = 2 - \frac{1}{1+x^2} + \frac{1}{\sqrt{1+x^2} - x} \left(\frac{x}{\sqrt{1+x^2}} - 1 \right)$$

$$\Rightarrow f'(x) = \frac{1+2x^2}{1+x^2} - \frac{1}{\sqrt{1+x^2}}$$

$$\Rightarrow f'(x) = \frac{1+2x^2}{1+x^2} - \frac{\sqrt{1+x^2}}{1+x^2}$$

$$\Rightarrow f'(x) = \frac{x^2 + \sqrt{1+x^2}\{\sqrt{1+x^2} - 1\}}{1+x^2} > 0 \text{ for all } x$$

Hence, $f(x)$ is an increasing function on $(-\infty, \infty)$ and in particular on $(0, \infty)$

7

(c)

We have,

$$f(x) = 3\cos^2 x + 4\sin^2 x + \cos \frac{x}{2} + \sin \frac{x}{2}$$

$$\Rightarrow f(x) = 4 - \cos^2 x + \cos \frac{x}{2} + \sin \frac{x}{2}$$



$$\begin{aligned} \Rightarrow f'(x) &= \sin 2x - \frac{1}{2} \left(\sin \frac{x}{2} - \cos \frac{x}{2} \right) \quad \dots(i) \\ \Rightarrow f'(x) &= 2 \sin x \cos x - \frac{1}{2} \left(\sin \frac{x}{2} - \cos \frac{x}{2} \right) \\ \Rightarrow f'(x) &= 2 \sin x \left(\sin^2 \frac{x}{2} - \cos^2 \frac{x}{2} \right) + \frac{1}{2} \left(\sin \frac{x}{2} - \cos \frac{x}{2} \right) \\ \Rightarrow f'(x) &= \left(\cos \frac{x}{2} - \sin \frac{x}{2} \right) \left\{ 2 \sin x \left(\cos \frac{x}{2} + \sin \frac{x}{2} \right) + \frac{1}{2} \right\} \\ \Rightarrow f'(x) &= \left(\cos \frac{x}{2} - \sin \frac{x}{2} \right) \left\{ 2\sqrt{2} \sin x \sin \left(\frac{x}{2} + \frac{\pi}{4} \right) + \frac{1}{2} \right\} \end{aligned}$$

For local maximum or minimum, we must have

$$f'(x) = 0$$

$$\Rightarrow \cos \frac{x}{2} - \sin \frac{x}{2} = 0$$

$$\Rightarrow \cos \frac{x}{2} = \sin \frac{x}{2} \Rightarrow \tan \frac{x}{2} = 1 \Rightarrow \frac{x}{2} = \frac{\pi}{4} \Rightarrow x = \frac{\pi}{2}$$

Now,

$$f''(x) = 2 \cos 2x - \frac{1}{4} \left(\cos \frac{x}{2} + \sin \frac{x}{2} \right) \quad [\text{Using (i)}]$$

$$\Rightarrow f'' \left(\frac{\pi}{2} \right) = 2 \cos \pi - \frac{1}{4} \left(\cos \frac{\pi}{4} + \sin \frac{\pi}{4} \right) = -2 - \frac{1}{2\sqrt{2}} < 0$$

Thus, $f(x)$ attains a local maximum at $x = \frac{\pi}{2}$

$$\text{Local maximum value} = f \left(\frac{\pi}{2} \right) = 4 + \frac{2}{\sqrt{2}} = 4 + \sqrt{2}$$

8

(c)

$$\therefore y = \left(\frac{c^6 - a^2 x^4}{b^2} \right)^{\frac{1}{4}}$$

$$\text{Let } f(x) = xy = \left(\frac{c^6 x^4 - a^2 x^8}{b^2} \right)^{\frac{1}{4}}$$

$$\Rightarrow f'(x) = \frac{1}{4} \left(\frac{c^6 x^4 - a^2 x^8}{b^2} \right)^{-3/4} \left(\frac{4x^3 c^6}{b^2} - \frac{8x^7 a^2}{b^2} \right)$$

$$\text{Put } f'(x) = 0$$

$$\Rightarrow x = \pm \frac{c^{3/2}}{2^{1/4} \sqrt{a}}$$

$$\therefore f \left(\frac{c^{3/2}}{c^{1/4} \sqrt{a}} \right) = \frac{c^3}{\sqrt{2ab}}$$

9

(a)

Let the radius of the circular wave ring by r cm at any time t . Then, $\frac{dr}{dt} = 30$ cm/sec (given)

Let A be the area of the enclosed ring. Then,

$$A = \pi r^2$$

$$\Rightarrow \frac{dA}{dt} = 2\pi r \frac{dr}{dt}$$

$$\Rightarrow \frac{dA}{dt} = 2\pi \times 50 \times \frac{30}{100} \text{ m}^2 \text{ sec} = 30\pi^2 \text{ m}^2 \text{ /sec}$$

10

(b)

We have,

$$x = t \cos t \text{ and } y = t \sin t$$

$$\therefore \frac{dx}{dt} = \cos t - t \sin t \text{ and } \frac{dy}{dx} = \sin t + t \cos t$$

At the origin, we have

$$x = 0, y = 0 \Rightarrow t \cos t = 0 \text{ and } t \sin t = 0 \Rightarrow t = 0$$



The slope of the tangent at $t = 0$ is

$$\frac{dy}{dx} = \left(\frac{\frac{dy}{dt}}{\frac{dx}{dt}} \right)_{t=0} = \left(\frac{\sin t + t \cos t}{\cos t - t \sin t} \right)_{t=0} = 0$$

So, the equation of the tangent at the origin is

$$t - 0 = 0(x - 0) \Rightarrow y = 0$$

11

(c)

Surface area of sphere $S = 4\pi r^2$ and $\frac{dr}{dt} = 2$

$$\therefore \frac{dS}{dt} = 4\pi \times 2r \frac{dr}{dt} = 8\pi r \times 2 = 16\pi r$$

$$\Rightarrow \frac{dS}{dt} \propto r$$

12

(c)

Let $f(x) = \left(\frac{1}{x}\right)^x = x^{-x} = e^{-x \log x}$. Then,

$$f'(x) = -\left(\frac{1}{x}\right)^x (\log x + 1) = -x^{-x}(\log x + 1)$$

Now,

$$f'(x) = 0$$

$$\Rightarrow -x^{-x}(\log x + 1) = 0$$

$$\Rightarrow \log x + 1 = 0 \Rightarrow \log x = -1 \Rightarrow x = e^{-1}$$

Clearly, $f''(x) < 0$ at $x = e^{-1}$

Hence, $f(x) = x^{-x}$ is maximum for $x = e^{-1}$. The maximum value is $e^{1/e}$

13

(c)

Given, $f(x) = 2x^3 - 21x^2 + 36x - 30$

$$\Rightarrow f'(x) = 6x^2 - 42x + 36$$

For maxima or minima, put $f'(x) = 0$

$$\Rightarrow 6x^2 - 42x + 36 = 0 \Rightarrow x = 6, 1$$

And $f''(x) = 12x - 42$

$$f''(1) = -30 \quad \text{and} \quad f''(6) = 30$$

Hence, $f(x)$ has maxima at $x = 1$ and minima at $x = 6$

14

(a)

Let l be the length of an edge and V be the volume of cube at any time t .

$$\therefore V = l^3$$

$$\therefore \frac{dV}{dt} = 3l^2 \frac{dl}{dt}$$

$$= 3 \times 5^2 \times 10 \text{ cm}^3/\text{s}$$

$$= 750 \text{ cm}^3/\text{s}$$

15

(c)

$$\text{We have, } \frac{dy}{dx} = \frac{-\sin \theta}{1 - \cos \theta}$$

Clearly, $\frac{dy}{dx} = 0$ for $\theta = (2k + 1)\pi$

So, the tangent is parallel to x -axis i.e. $y = 0$

16

(b)

We have,

$$5x^5 - 10x^3 + x + 2y + 6 = 0 \quad \dots(i)$$

Differentiating with respect to x , we get

$$25x^4 - 30x^2 + 1 + 2 \frac{dy}{dx} = 0$$

$$\Rightarrow \frac{dy}{dx} = -\frac{1}{2}(25x^4 - 30x^2 + 1)$$



$$\Rightarrow \left(\frac{dy}{dx}\right)_{(0,-3)} = -\frac{1}{2}$$

The equation of the normal at $(0, -3)$ is

$$y + 3 = 2(x - 0) \Rightarrow 2x - y - 3 = 0$$

Solving (i) and (ii), we obtain the coordinates of their points of intersection as $P(0, -3)$, $(1, -1)$ and $(-1, -5)$

Hence, the normal at $P(0, -3)$ meets the curve again at $(1, -1)$ and $(-1, -5)$

17

(b)

We have,

$$\frac{dx}{d\theta} = a(-\sin\theta + \sin\theta + \theta \cos\theta) = a\theta \cos\theta$$

$$\text{and, } \frac{dy}{d\theta} = a(\cos\theta - \cos\theta + \theta \sin\theta) = a\theta \sin\theta$$

$$\therefore \frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \tan\theta \Rightarrow -\frac{1}{\frac{dy}{dx}} = -\cot\theta$$

Hence, the slope of the normal varies as θ

The equation of the normal at any point is

$$y - a(\sin\theta - \theta \cos\theta) = -\cot\theta \{x - a(\cos\theta + \theta \sin\theta)\}$$

$$\Rightarrow x \cos\theta + y \sin\theta = a$$

Clearly, it is a line at a constant distance $|a|$ from the origin

18

(d)

We have,

$$f(x) = \begin{cases} 3x^2 + 12x - 1, & -1 \leq x \leq 2 \\ 37 - x, & 2 < x \leq 3 \end{cases}$$

$$\text{Clearly, } \lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x) = f(2)$$

So, $f(x)$ is continuous at $x = 2$

Hence, it is continuous on $[-1, 3]$

Thus, option (b) is correct

We find that

$$f'(x) = 6x + 12 > 0 \text{ for all } x \in [-1, 2]$$

$\Rightarrow f(x)$ is increasing on $[-1, 2]$

Thus, option (a) is correct

Also,

$$f'(x) < 0 \text{ for all } x \in (2, 3]$$

$\Rightarrow f(x)$ is decreasing on $(2, 3]$

Hence, $f(x)$ attains the maximum value at $x = 2$

So, option (c) is correct

19

(b)

$$\text{Given, } f(x) = x^3 - 3x^2 + 2x$$

$$\Rightarrow f'(x) = 3x^2 - 6x + 2$$

$$\text{Now, } f(a) = f(0) = 0$$

$$\text{And } f(b) = f\left(\frac{1}{2}\right)$$

$$= \frac{1}{2} \left(\frac{1}{2} - 1\right) \left(\frac{1}{2} - 2\right) = \frac{3}{8}$$

By Lagrange's Mean Value Theorem

$$\frac{f(b) - f(a)}{b - a} = f'(c)$$

$$\Rightarrow \frac{\frac{3}{8} - 0}{\frac{1}{2} - 0} = 3c^2 - 6c + 2$$

$$\Rightarrow 12c^2 - 24c + 5 = 0$$



This is a quadratic equation in c.

$$c = \frac{24 \pm \sqrt{576 - 240}}{24}$$

$$= 1 \pm \frac{\sqrt{21}}{6}$$

But c lies between 0 to $\frac{1}{2}$

$$\therefore \text{we take, } c = 1 - \frac{\sqrt{21}}{6}$$

20

(a)

Since, $f(x) = kx - \sin x$ is monotonically increasing for all $x \in R$. Therefore,

$$f'(x) > 0 \text{ for all } x \in R$$

$$\Rightarrow K - \cos x > 0$$

$$\Rightarrow K > \cos x$$

$$\Rightarrow K > 1 \text{ [}\because \text{maximum value of } \cos x \text{ is } 1]$$

ANSWER-KEY

Q.	1	2	3	4	5	6	7	8	9	10
A.	B	A	C	C	C	A	C	C	A	B
Q.	11	12	13	14	15	16	17	18	19	20
A.	C	C	C	A	C	B	B	D	B	A

**SMARTLEARN
COACHING**