





CLASS : XIth DATE : SUBJECT : CHEMISTRY DPP No. : 2

## **Topic :-SOLUTIONS**

- 1. Which is not applicable to distribution law?
  - a) Parke's process
  - b) Solvent extraction
  - c) Pattinson's process
  - d) Partition chromatography
- 2. Which of the following is the expression of Raoult's law?
  - $(p = vapour pressure of pure solvent, <math>p_s = vapour pressure of the solution)$

a) 
$$\frac{p - p_s}{p} = \frac{n}{n + N}$$
 b)  $\frac{p_s - p}{p} = \frac{N}{N + n}$  c)  $\frac{p - p_s}{p_s} = \frac{N}{N - n}$  d)  $\frac{p_s - p}{p_s} = \frac{N - n}{N}$ 

- 3. For determination of molar mass of colloids, polymers and protein, which property is used ?
  - a) Diffusion pressure c) Osmotic pressure
- b) Atmospheric pressure d) Turgor pressure
- 4.3.0 molal NaOH solution has a density of 1.110 g/mL. The molarity of the solution is<br/>a) 3.9732b) 2.9732c) 1.9732d) 0.9732
- Sodium sulphate is soluble in water, while barium sulphate is sparingly soluble because :
   a) The hydration energy of sodium sulphate is more than its lattice energy
  - b) The lattice energy of barium sulphate is less than the hydration energy
  - c) The lattice energy has no role to play in solubility
  - d) The hydration energy of sodium sulphate is less than its lattice energy
- 6. Distribution law is applicable when :
  - a) Temperature remains constant
  - b) Dilute solutions are employed
  - c) The two solvents are mutually insoluble
  - d) All are correct
- 7.10 cm³ of 0.1 N monobasic acid requires 15 cm³ of sodium hydroxide solution whose normality is<br/>a) 1.5 Nc) 0.066 Nd) 0.66 N
- 8. Density of a 2.05 M solution of acetic acid in water is 1.02 g/mL. The molality of the solution is a) 1.14 mol  $kg^{-1}$  b) 3.28 mol  $kg^{-1}$  c) 2.28 mol  $kg^{-1}$  d) 0.44 mol  $kg^{-1}$

9. x gram of water is mixed in 69 g of ethanol. Mole fraction of ethanol in the resultant solution is 0.6. What is the value of x in grams?
a) 54 b) 36 c) 180 d) 18

10. Dissolution of a solute is an exothermic process if :a) Hydration energy > lattice energy

| 5        |   |  | Sma  | rt DPPs   |
|----------|---|--|--|---|
| COACHING |   |  |  |   |
|          | <ul> <li>b) Hydration energy &lt;</li> <li>c) Hydration energy =</li> <li>d) None of the above</li> </ul>   |  |  |   |
| 11.      | Molarity is expressed a<br>a) L/mol   | s<br>b) Mol/L  | c) Mol/1000 g  | d) g/L  |
| 12.      | The amount of anhydro<br>a) 6.0 g   | ous Na <sub>2</sub> CO <sub>3</sub> present in 250<br>b) 6.625 g | mL of 0.25 M solution is<br>c) 66.25 g                         | d) 6.225 g  |
| 13.      | Which of the following<br>a) HCl  | compounds correspond to<br>b) MgSO <sub>4</sub>                  | o maximum van'thoff facto<br>c) K <sub>2</sub> SO <sub>4</sub> | r for dilute solution?<br>d) K <sub>4</sub> Fe(CN) <sub>6</sub> |
| 14.      | Solute <i>A</i> is a ternary electrolyte and solute <i>B</i> is non-electrolyte. If 0.1 <i>M</i> solution of solute <i>B</i> produces an osmotic pressure of 2 <i>P</i> , then 0.05 <i>M</i> solution of <i>A</i> at the same temperature will produce an osmotic pressure equal to : |  |  |   |
|          | a) P  | b) 1.5 <i>P</i>  | c) 2 <i>P</i>  | d) 3 <i>P</i>   |
| 15.      | A solution of sucrose (molar mass $342 \text{ g mol}^{-1}$ ) has been produced by dissolving 68.5 g sucrose in 1000 g water. The freezing point of the solution obtained will be : ( $K_f$ for H <sub>2</sub> O = 1.86 Kkg mol <sup>-1</sup> )  |  |  |   |
|          | a) –0.372°C   | b) -0.520°C  | c) + 0.372°C   | d) -0.570°C   |
| 16.      | A mixture of ethyl alcohol and propylalcohol has a vapour pressure of 290 mm at 300 K. the vapour pressure of propylalcohol is 200 mm. If the mole fraction of ethylalcohol is 0.6, its vapour pressure (in mm) at the same temperature will be                                       |  |  |   |
|          | a) 350  | b) 300   | c) 700   | d) 360  |
| 17.      | How many grams of sulphuric acid is to be dissolved to prepare 200 mL aqueous solution having concentration of $[H_3O^+]$ ions 1 M at 25°C temperature.<br>$[H = 1, O = 16, S = 32 \text{ g.mol}^{-1}]$   |  |  |   |
|          | a) 4.9g   | b) 19.6g   | c) 9.8g  | d) 0.98g  |
| 18.      | 3. What is the molarity of $H_2SO_4$ solution if 25 mL is exactly neutralised with 32.63 mL of 0.164 M  |  |  |   |
|          | NaOH?<br>a) 0.107 M   | b) 0.126 M   | c) 0.214 M   | d) -0.428 M   |
| 19.      | What is the molality $-10^{\circ}$ C?.( $K_f$ for water =   |  | wt. = 416) in aqueous  | solution which freezes at                                       |
|          | a) 3.540  | b) 4.567   | c) 5.376   | d) 6.315  |
| 20.      | The solubility order for<br>a) $NH_3 > CO_2 > O_2 > D_2$<br>b) $H_2 > O_2 > NH_3 > C$<br>c) $CO_2 > NH_3 > O_2 > D_2$<br>d) $O_2 > H_2 > NH_3 > C$  | H <sub>2</sub><br>O <sub>2</sub><br>N <sub>2</sub>               |  |   |