

DPP

DAILY PRACTICE PROBLEMS

CLASS : XIIth
DATE :

SUBJECT : MATHS
DPP NO. : 3

Topic :-MATRICES

- If A and B are matrices such that AB and $A + B$ both are defined, then
 - A and B can be any two matrices
 - A and B are square matrices not necessarily of the same order
 - A, B are square matrices of the same order
 - Number of columns of A is same as the number of rows of B
- Let a, b, c be any real numbers. Suppose that there are real numbers x, y, z not all zero such that $x = cy + bz, y = az + cx,$ and $z = bx + ay$ have non-zero solution. Then, $a^2 + b^2 + c^2 + 2abc$ is equal to
 - 1
 - 2
 - 1
 - 0
- If I_n is the identity matrix of order n , then rank of I_n is
 - 1
 - n
 - 0
 - None of these
- If the matrix $A = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & \lambda \end{bmatrix}$ is singular, then λ is equal to
 - 3
 - 4
 - 2
 - 5
- If $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$, then $I + A + A^2 + A^3 + \dots \infty$ equals to
 - $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
 - $\begin{bmatrix} -1 & -2 \\ -3 & -4 \end{bmatrix}$
 - $\begin{bmatrix} 1/2 & -1/3 \\ -1/2 & 0 \end{bmatrix}$
 - $\begin{bmatrix} -1/4 & 1/3 \\ 1/2 & 0 \end{bmatrix}$
- If A is a non-singular square matrix of order n , then the rank of A is
 - Equal to n
 - Less than n
 - Greater than n
 - None of these
- If $A = \begin{bmatrix} 1 & -2 \\ 4 & 5 \end{bmatrix}$ and $f(t) = t^2 - 3t + 7$, then $f(A) + \begin{bmatrix} 3 & 6 \\ -12 & -9 \end{bmatrix}$ is equal to
 - $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
 - $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$
 - $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$
 - $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$
- The system of linear equations

$$\begin{aligned} x + y + z &= 2 \\ 2x + y - z &= 3 \\ 3x + 2y + kz &= 4 \end{aligned}$$
 has a unique solution if
 - $k \neq 0$
 - $-1 < k < 1$
 - $-2 < k < 2$
 - $k = 0$
- The number of solutions of the system of equations

$$\begin{aligned} 2x + y - z &= 7, \\ x - 3y + 2z &= 1, \\ x + 4y - 3z &= 5 \end{aligned}$$
 is
 - 0
 - 1
 - 2
 - 3
- If $X = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$, the value of X^n is equal to



- a) $\begin{bmatrix} 3n & -4n \\ n & -n \end{bmatrix}$ b) $\begin{bmatrix} 2+n & 5-n \\ n & -n \end{bmatrix}$ c) $\begin{bmatrix} 3^n & (-4)^n \\ 1^n & (-1)^n \end{bmatrix}$ d) None of these
11. If I_3 is the identity matrix of order 3, then $(I_3)^{-1} =$
a) 0 b) $3 I_3$ c) I_3 d) Not necessarily exists
12. If $A = [a_{ij}]$ is a square matrix of order $n \times n$ and k is a scalar, then $|kA| =$
a) $k^n |A|$ b) $k|A|$ c) $k^{n-1}|A|$ d) None of these
13. If $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ a & b & -1 \end{bmatrix}$, then A^2 is equal to
a) Null matrix b) Unit matrix c) $-A$ d) A
14. If $A = \begin{bmatrix} \alpha & 0 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 \\ 5 & 1 \end{bmatrix}$, then value of α for which $A^2 = B$ is
a) 1 b) -1 c) 4 d) No real values
15. If A is a square matrix such that $A (\text{adj } A) = \begin{bmatrix} 4 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 4 \end{bmatrix}$, then $|\text{adj } A| =$
a) 4 b) 16 c) 64 d) 256
16. If ω is a complex cube root of unity and $A = \begin{bmatrix} \omega & 0 \\ 0 & \omega \end{bmatrix}$, then A^{50} is
a) $\omega^2 A$ b) ωA c) A d) 0
17. If $A = \begin{bmatrix} 1 & 2 & x \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & -2 & y \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ and $AB = I_3$, then $x + y$ equals
a) 0 b) -1 c) 2 d) None of these
18. The adjoint of the matrix $\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$ is
a) $\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ b) $\begin{bmatrix} \sin \theta & \cos \theta \\ \cos \theta & \sin \theta \end{bmatrix}$ c) $\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$ d) $\begin{bmatrix} -\sin \theta & \cos \theta \\ \cos \theta & \sin \theta \end{bmatrix}$
19. The inverse matrix of $A = \begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{bmatrix}$ is
a) $\begin{bmatrix} \frac{1}{2} & -\frac{11}{22} \\ -4 & 3 & -1 \\ \frac{5}{2} & -\frac{31}{22} \end{bmatrix}$ b) $\begin{bmatrix} \frac{1}{2} & -4 & \frac{5}{2} \\ 1 & -6 & 3 \\ 1 & 2 & -1 \end{bmatrix}$ c) $\frac{1}{2} \begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \\ 4 & 2 & 3 \end{bmatrix}$ d) $\frac{1}{2} \begin{bmatrix} 1 & -1 & -1 \\ -8 & 6 & -2 \\ 5 & -3 & 1 \end{bmatrix}$
20. If $f(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$, then $\{f(\theta)^{-1}\}$ is equal to
a) $f(-\theta)$ b) $f(\theta)^{-1}$ c) $f(2\theta)$ d) None of these



SMARTLEARN
COACHING