

DPP

DAILY PRACTICE PROBLEMS

CLASS : XIIth
DATE :

SUBJECT : MATHS
DPP NO. : 3

Topic :-INTEGRALS

- If $f(x) = \lim_{n \rightarrow \infty} [2x + 4x^3 + \dots + 2nx^{2n-1}] (0 < x < 1)$, then $\int f(x) dx$ is equal to
 a) $-\sqrt{1-x^2}$ b) $\frac{1}{\sqrt{1-x^2}}$ c) $\frac{1}{x^2-1}$ d) $\frac{1}{1-x^2}$
- $\int \frac{dx}{\sin x - \cos x + \sqrt{2}}$ equals
 a) $-\frac{1}{\sqrt{2}} \tan\left(\frac{x}{2} + \frac{\pi}{8}\right) + c$ b) $\frac{1}{\sqrt{2}} \tan\left(\frac{x}{2} + \frac{\pi}{8}\right) + c$ c) $\frac{1}{\sqrt{2}} \cot\left(\frac{x}{2} + \frac{\pi}{8}\right) + c$ d) $-\frac{1}{\sqrt{2}} \cot\left(\frac{x}{2} + \frac{\pi}{8}\right) + c$
- $\int_0^{\pi/2} \frac{\cos x}{1+\sin x} dx$ is equal to
 a) $\log 2$ b) $2 \log 2$ c) $(\log 2)^2$ d) $\frac{1}{2} \log 2$
- The integral $\int_0^1 \frac{2 \sin^{-1} x}{x} dx$ equals
 a) $\int_0^{\pi/6} \frac{x}{\tan x} dx$ b) $\int_0^{\pi/6} \frac{2x}{\tan x} dx$ c) $\int_0^{\pi/2} \frac{2x}{\tan x} dx$ d) $\int_0^{\pi/6} \frac{x}{\sin x} dx$
- If $\int_2^e \left(\frac{1}{\log x} - \frac{1}{(\log x)^2} \right) dx = a + \frac{b}{\log 2}$, then
 a) $a = e, b = -2$ b) $a = e, b = 2$ c) $a = -e, b = 2$ d) None of these
- The value of $\int_0^8 |x - 5| dx$ is
 a) 17 b) 12 c) 9 d) 18
- $\int_0^1 \frac{x dx}{[x + \sqrt{1-x^2}] \sqrt{1-x^2}}$ is equal to
 a) 0 b) 1 c) $\frac{\pi}{4}$ d) $\frac{\pi^2}{2}$
- $\int_0^1 \cot^{-1}(1 - x + x^2) dx$ is equal to
 a) $\pi - \log 2$ b) $\pi + \log 2$ c) $\frac{\pi}{2} + \log 2$ d) $\frac{\pi}{2} - \log 2$
- $\int_8^{15} \frac{dx}{(x-3)\sqrt{x+1}}$ is equal to
 a) $\frac{1}{2} \log \frac{5}{3}$ b) $\frac{1}{3} \log \frac{5}{3}$ c) $\frac{1}{2} \log \frac{3}{5}$ d) $\frac{1}{5} \log \frac{3}{5}$
- $\int \{1 + 2 \tan x (\tan x + \sec x)\}^{1/2} dx$ is equal to
 a) $\log \sec x (\sec x - \tan x) + C$
 b) $\log \operatorname{cosec}(\sec x + \tan x) + C$



- c) $\log \sec x (\sec x + \tan x + C)$
d) $\log(\sec x + \tan x) + C$
11. If $I_1 = \int_0^\infty \frac{1}{1+x^4} dx$ and $I_2 = \int_0^\infty \frac{x^2}{1+x^4} dx$. Then $\frac{I_1}{I_2} =$
a) 1 b) 2 c) 1/2 d) None of these
12. $\int \frac{\sin x dx}{3+4 \cos^2 x}$ is equal to
a) $\log(3 + 4 \cos^2 x) + c$ b) $\frac{1}{2\sqrt{3}} \tan^{-1} \left(\frac{\cos x}{\sqrt{3}} \right) + c$
c) $-\frac{1}{2\sqrt{3}} \tan^{-1} \left(\frac{2\cos x}{\sqrt{3}} \right) + c$ d) $\frac{1}{2\sqrt{3}} \tan^{-1} \left(\frac{2\cos x}{\sqrt{3}} \right) + c$
13. For any integer n , the integral $\int_0^\pi e^{\cos^2 x} \cos^3(2n + 1)x dx$ has the value
a) π b) 1 c) 0 d) None of these
14. If $\frac{d}{dx} \{f(x) = \frac{1}{1+x^2}\}$, then $\frac{d}{dx} \{f(x^3)\}$ is
a) $\frac{3x}{1+x^3}$ b) $\frac{3x^2}{1+x^6}$ c) $\frac{-6x^5}{(1+x^6)^2}$ d) $\frac{-6x^5}{1+x^6}$
15. $\int_0^\pi [\cot x] dx$, $[.]$ denotes the greatest integer function, is equal to
a) $\frac{\pi}{2}$ b) 1 c) -1 d) $-\frac{\pi}{2}$
16. $\int_{-3}^2 \{|x + 1| + |x + 2| + |x - 1|\} dx$ is equal to
a) $\frac{31}{2}$ b) $\frac{35}{2}$ c) $\frac{47}{2}$ d) $\frac{39}{2}$
17. $\int_0^3 |x^3 + x^2 + 3x| dx$ is equal to
a) $\frac{171}{2}$ b) $\frac{171}{4}$ c) $\frac{170}{4}$ d) $\frac{170}{3}$
18. $\int \frac{dx}{\sin x \cos x}$ is equal to
a) $\log |\sin x| + c$ b) $\log |\tan x| + c$ c) $\log |\sec x| + c$ d) None of these
19. $\int_0^{2n\pi} \left\{ |\sin x| - \left| \frac{1}{2} \sin x \right| \right\} dx$ equals
a) nb b) $2nc$ c) $-2nd$ d) None of these
20. If $\int_a^b x^3 dx = 0$ and if $\int_a^b x^2 dx = \frac{2}{3}$, then the values of a and b are respectively
a) 1,1 b) -1, -1 c) 1, -1 d) -1,1