



## REDOX REACTION

### 1 BASIC CONCEPT

In the previous chapter, we have studied neutralization reactions in which  $H^+$  ions from an acid ( $HCl$ ,  $H_2SO_4$ ,  $CH_3COOH$  etc.) combine with  $OH^-$  ions of a base ( $NaOH$ ,  $Ca(OH)_2$ ,  $NH_4OH$  etc.) to form a salt and weakly ionized molecules of water. In this chapter, we shall discuss another important class of reactions called reduction–oxidation or simply redox reactions. All these reactions are always accompanied by energy changes in the form of heat, light or electricity.


A large number of chemical and biological reactions fall in this category. Burning of different types of fuels for obtaining energy for domestic, transport or industrial purpose, electrochemical processes like manufacturing of caustic soda, digestion of food in animals, photosynthesis by plants, corrosion of metals and operations of dry and wet batteries are diverse examples of redox reactions.

### 2 OXIDATION AND REDUCTION

In redox reactions one reactant called **reductant or reducing agent** is oxidised while the other reactant called **oxidant or oxidising agent** is reduced.

#### 2.1 CLASSICAL CONCEPT OF OXIDATION AND REDUCTION


Originally, **oxidation** of a substance was defined as the addition of oxygen or any other electronegative element to it. Due to the presence of oxygen in the atmosphere, many elements combine with it. In fact the most common occurrence of elements on the earth is in the form of their oxides. **Later on, removal of hydrogen or any other electropositive element from a substance was also considered as oxidation.**




Magnesium is oxidised by addition of oxygen to it.



Hydrogen sulphide is oxidised by the removal of hydrogen from it.






Magnesium is oxidised by the addition of fluorine to it.



Magnesium is oxidised by the addition of sulphur to it.

Similarly reduction of a substance was initially introduced for the processes in which metal oxides were converted into metals by their reactions with hydrogen, carbon or other metals. Carbon monoxide is also used in the reduction of ferric oxide to iron.



Here, in all the three cases mentioned above metal oxide is reduced by hydrogen, carbon or carbon monoxide.

#### 2.2 MODERN CONCEPT OF OXIDATION AND REDUCTION


The general definition of oxidation is **loss of electrons**. In the above examples of oxidation magnesium is converted into  $Mg^{2+}$  ion and in the process it loses two electrons.

Similarly, reduction in general means **acceptance of electrons**. When  $CuO$  reacts with hydrogen it gets converted into copper. In copper oxide, each  $Cu^{2+}$  ion accepts two electrons and gets reduced to copper.




## 2.3 COMPLIMENTARY NATURE OF OXIDATION AND REDUCTION REACTIONS

Whenever any substance is oxidised, another substance is always reduced at the same time and vice-versa. In other words, oxidation and reduction reactions are complimentary. This is illustrated by the following examples




Here  $\text{H}_2\text{S}$  is oxidised to  $\text{S}$  while  $\text{Cl}_2$  is reduced to  $\text{HCl}$ .




Here,  $\text{HCl}$  is oxidised to  $\text{Cl}_2$  while  $\text{MnO}_2$  is reduced to  $\text{MnCl}_2$ .


The occurrence of electron transfer is more apparent in some redox reactions than others. When metallic zinc is added to a solution containing copper(II) sulphate, zinc reduces  $\text{Cu}^{2+}$  by donating two electrons to it.





In the process, solution loses the blue colour that characterizes the presence of  $\text{Cu}^{2+}$  ions. The oxidation and reduction reactions are



Similarly, metallic copper reduces silver ions in a solution of silver nitrate ( $\text{AgNO}_3$ )





The oxidation and reduction reactions are



### PRACTICE PROBLEM

PP1. Identify the oxidant and the reductant in the following reactions.



## 3 THE OXIDATION STATE OF AN ELEMENT

To describe the changes that occur in oxidation–reduction reactions and to write the correctly balanced equations for such reactions, it is helpful to know the concept of the oxidation state (or oxidation number) of an atom.

For monoatomic ions, **the oxidation state is simply the charge on the ion**. For example, oxidation states of  $\text{Na}^+$ ,  $\text{Mg}^{2+}$ ,  $\text{Cl}^-$  and  $\text{N}^{3-}$  are +1, +2, –1 and –3 respectively. For covalently bonded substances, the charge on an atom would be so small that it is impossible to calculate the exact charge on each atom in the molecule. Thus, oxidation state of an atom in a covalently bonded molecule is defined as **“the charge an atom would possess if all the bonds associated with that atom are broken heterolytically considering them to be completely ionic”**. For example, in  $\text{NH}_3$  there are three N–H bonds. If we consider all three N–H bonds to be ionic, then each H will possess a charge of +1 while N acquires a charge of –3 (because N is more electronegative than H). In  $\text{H}_2\text{O}_2$ , there are two O–H bonds and one O–O bond as evident from the structural formula H–O–O–H. Assuming each O–H bond to be ionic, each H atom would possess +1 charge while each O would possess –1 charge. The O–O bond is not ionic as the bond is between atoms of similar electronegativity. Thus oxidation state of each O-atom in  $\text{H}_2\text{O}_2$  is –1.

It is not necessary to know the structure of a molecule in order to calculate the oxidation state of an atom in a molecule. The following set of rules are used to assign oxidation state to each atom in an ion or molecule.

1. Each pure element has an oxidation state of zero. This is true whether the element is a monoatomic gas, a metallic solid or a polyatomic molecule. Thus,  $\text{Fe}(\text{s})$ ,  $\text{N}_2(\text{g})$ ,  $\text{P}_4(\text{s})$  and  $\text{S}_8(\text{s})$  are all in the zero oxidation state.



- In monoatomic ions, the oxidation state of the element is equal to the charge on the ion. Thus, oxidation state of all alkali metals is +1 and that of alkaline earth metals is +2 in all their compounds. In ionic solid  $K_2S$ , the oxidation states of potassium and sulfur are +1 and  $-2$  respectively. In  $AgCl$ , the oxidation state of silver and chlorine are +1 and  $-1$  respectively.
- The oxidation state of hydrogen in any molecule in which it is combined with another element is +1, except in the metallic hydrides such as  $LiH$  or  $CaH_2$ , where the oxidation state of hydrogen is  $-1$ .
- The oxidation state of oxygen in any molecule or ion in which it is combined with another element is  $-2$ , except in the peroxides, the superoxides and in  $OF_2$ . The peroxides ( $H_2O_2$ ,  $Na_2O_2$ ,  $BaO_2$  etc.) are compounds in which there is an  $O-O$  covalent bond and the peroxide ion is  $O_2^{2-}$ . The oxidation state of oxygen in peroxides is  $-1$ . The superoxides (like  $KO_2$ ,  $CsO_2$  etc.) are ionic compounds involving the superoxide ion,  $O_2^-$ . The oxidation state of oxygen is  $-\frac{1}{2}$  in superoxide ion. In  $OF_2$ , the oxidation state of oxygen is +2 since fluorine is more electronegative than oxygen.
- The oxidation state of all the halogens is  $-1$  in all their compounds except where they are combined with an element of higher electronegativity or in interhalogen compounds.. Oxidation state of fluorine is always  $-1$ , since it is the most electronegative element.
- In covalent compounds not involving hydrogen or oxygen, the more electronegative element is assigned negative oxidation state while a less electronegative element is assigned a positive oxidation state.
- The algebraic sum of the oxidation numbers of all the atoms combined in a molecule or complex ion must equal the net charge on the molecule or ion.

$Fe_3O_4$  is made up of equimolar quantity of  $FeO$  and  $Fe_2O_3$ .


## 4 TYPES OF REDOX REACTION

There are four general types of redox reactions, namely

- Combination reactions.
- Decomposition reactions.
- Displacement reactions.
- Disproportionation reactions.


### 4.1 COMBINATION REACTIONS

A combination reaction may be represented by




If either A or B is an element, then the reaction is usually redox in nature. **Combination reactions** are reactions in which two or more substances combine to form a single product.

For example,



### 4.2 DECOMPOSITION REACTIONS


Decomposition reactions are the opposite of combination reactions. Specifically, a **decomposition reaction** is the breakdown of a compound into two or more components.



Note that we show oxidation states only for the elements that are oxidized or reduced.

## 4.3 DISPLACEMENT REACTIONS



In a *displacement reaction*, an ion (or atom) in a compound is replaced by an ion (or atom) of another element:



Most displacement reactions fit into one of three subcategories: hydrogen displacement, metal displacement or halogen displacement.

### 4.3.1. Hydrogen Displacement



All alkali metals and some alkaline earth metals (Ca, Sr and Ba), which are the most reactive of the metallic elements, will displace hydrogen from cold water.



Less reactive metals, such as aluminium and iron, react with steam to give hydrogen gas:



Many metals, including those that do not react with water, are capable of displacing hydrogen from acids. For example, zinc (Zn) and magnesium (Mg) do not react with water but do react with hydrochloric acid, as follows:



### 4.3.2. Metal Displacement

A metal in a compound can be displaced by another metal in the elemental state.

In addition to the two examples given in section 2.3 some more examples are given as follows




### 4.3.3. Halogen Displacement

Another activity series summarizes the halogens behaviour in halogen displacement reactions:




The power of these elements as oxidizing agents decreases as we move down the group VII A from fluorine to iodine. So molecular fluorine can displace chloride, bromide and iodide ions in solution. In fact, molecular fluorine is so reactive that it also attacks water. Thus, these reactions cannot be carried out in aqueous solutions. On the other hand molecular chlorine can displace bromide and iodide ions in aqueous solution. The displacement equations are



## 4.4 DISPROPORTIONATION REACTIONS

A special type of redox reaction is the *disproportionation reaction*. In this type of reactions, an element in one oxidation state is simultaneously oxidized and reduced. One reactant in a disproportionation reaction always contains an element that shows increase in its oxidation state in one product and decrease in its oxidation state in another product. The element itself is in an intermediate oxidation state, i.e., both higher and lower oxidation states of the element exist on the product side. The decomposition of hydrogen peroxide is an example of a disproportionation reaction





Redox reactions involve both oxidation and reduction simultaneously. Oxidation means loss of electrons and reduction means gain of electrons. Thus redox reactions involve electron transfer and the number of electrons lost is same as the number of electrons gained during the reaction. This aspect of redox reaction can serve as the basis of a pattern for balancing redox reactions.

There are two common and useful methods to balance redox reactions. These are

- Oxidation state method and
- Ion–electron method.

## 5.1 BALANCING OF REDOX REACTIONS BY OXIDATION STATE METHOD

For balancing a redox reaction by oxidation state method, follow the order of steps as listed below. In balancing some reactions, all the steps may not be required.

- For each redox reaction, deduce the oxidation state of the elements that are undergoing oxidation and reduction.
- Separate the reactants and products into two half–reactions involving the elements alone that show change in their oxidation states. Write the skeletal equations for each half–reaction.
- For each half–reaction, first balance the number of atoms of the element undergoing change in oxidation state.
- Now find the total change in oxidation state by determining the change per atom and multiplying it by the total number of atoms that undergo change in oxidation state. Also, decide whether electrons are lost or gained. An increase in oxidation state means loss of electrons and a decrease in oxidation state means gain of electrons.
- Add the electrons lost or gained to the half equation. Lost electrons are placed on the product side and gained electrons are kept on the reactant side.
- Now add both the half reactions after multiplying by suitable integers to make the number of electrons lost and gained same.
- Transfer the coefficients of each reactant and product to the main skeletal equation.
- If the coefficients developed are not correct, then change them by inspection method. Such coefficient changes are required when an element from a compound goes in 2 different compounds, one with the same oxidation state and the other with different oxidation state.
- Count the charges on both sides of the equation and balance the charges in the equation by adding requisite  $H^+$  or  $OH^-$  to the required side. If the reaction occurs in acidic solution, use  $H^+$  and if it occurs in basic solution, use  $OH^-$ . If the reaction occurs in neutral solution, use  $H^+$  or  $OH^-$  on any of the side as needed i.e. in a neutral solution, if negative charges are needed for balancing, use  $OH^-$  and if positive charges are needed, use  $H^+$ .
- Balance the hydrogens and oxygens by adding the appropriate number of  $H_2O$  molecules on the required side.

## 5.2 BALANCING REDOX REACTIONS BY ION–ELECTRON METHOD

This method of balancing redox reaction involves following steps.

- For each redox reaction, deduce the oxidation state of the elements that are undergoing oxidation and reduction.
- Separate the reactants and products into two half–reactions involving the elements that show change in their oxidation states. Write the skeleton equations for each half–reaction.
- Balance each half–reaction separately involving the given steps.
  - First balance the atoms of the element undergoing oxidation or reduction.
  - Then balance atoms of the elements other than hydrogen and oxygen.
  - For balancing oxygen atoms in acidic or neutral medium, add suitable number of  $H_2O$  molecules to the side deficient in O while in alkaline medium, add equal number of  $H_2O$  molecules as the excess of O on the side having excess of O atoms and add double the number of  $OH^-$  ions on the opposite side of the equation.
  - In order to balance the hydrogen atoms in acidic or neutral medium, add required number of  $H^+$  to the side deficient in H while in alkaline medium, add equal number of  $OH^-$  ions as the excess number of H atoms on the side having excess

H and add equal number of  $\text{H}_2\text{O}$  molecules on the opposite side of the equation.

- (iv) Multiply each half-reaction by suitable integer to make the number of electrons lost and gained same and add both the half-equations to get a completely balanced reaction.

Mostly, the medium in which a redox reaction is to be balanced is given in the problem but if the problem does not state the medium explicitly, then the medium is decided by looking at the reactants or products. If an acid or base is one of the reactants or products, then that reactant or product decides the medium. If ammonia is present, the solution would be basic, and if ammonium chloride is present, it would be acidic. If metals which form insoluble hydroxides are given in their ionic form in a redox reaction, the solution is acidic.

Although, both the oxidation state method and ion-electron method lead to the correct form of the balanced redox reaction but ion-electron method is considered to be superior to the oxidation state method due to following advantages:

- In ion-electron method, all reactants and products are completely balanced (i.e. coefficient is developed for each one of them) whether they participate in the redox change or not, while in oxidation state method, the balancing of coefficients is developed only for the species involved in the redox change.
- The balancing coefficients developed in ion-electron method are always correct and need no amendment while such coefficients are to be changed sometimes in oxidation number method.

## 6 ELECTROCHEMICAL CELLS

Let us take a zinc rod and immerse it in a solution of  $\text{CuSO}_4$  taken in a beaker as shown in the figure 1. After some time, we see that the Zn rod starts dissolving and Cu starts depositing on its surface. This happens because Zn gets oxidised to  $\text{Zn}^{2+}$  ions, which passes into the solution and 2 electrons remain on the Zn rod for each  $\text{Zn}^{2+}$  ion going in solution. Then,  $\text{Cu}^{2+}$  ion from the solution takes up these 2 electrons to form Cu, which deposits on the Zn rod.




Figure 1



In this reaction, Zn acts as reducing agent, which is able to reduce  $\text{Cu}^{2+}$  to Cu by transferring 2 electrons and  $\text{Cu}^{2+}$  ion acts as an oxidising agent, which oxidises Zn to  $\text{Zn}^{2+}$  ions and itself gets reduced to Cu. In this case, there is a direct transfer of electrons from Zn rod to  $\text{Cu}^{2+}$  ion and some heat is also evolved.

The above experiment is slightly modified in a manner that for the same redox reaction, transfer of electrons takes place indirectly and the heat of the reaction is converted into electrical energy. This necessitates the separation of Zn rod from  $\text{CuSO}_4$  solution. Such cells in which the oxidised and reduced species are separated and connected through electrical wires are called electrochemical cells.

**Electrochemical cells are the cells in which chemical energy is transformed into electrical energy. This means that chemical reactions produce electric current.**

An electrochemical cell consists of two half-cells or electrodes. The electrodes are metallic conductors dipped in an electrolyte, which is an ionic conductor dissolved in water. A conducting rod and a suitable electrolyte comprise an electrode or half-cell compartment. The two electrodes may share the same electrolyte or have different electrolytes.

In a given electrochemical cell, combination of any of the two electrodes can be used. The cell may even contain same type of electrodes with different concentration of electrolytes. When an 'inert metal' is part of the electrode, it acts as a source of electrons, but does not take part in the reaction. If the electrolytes are different, the two compartments may be joined by a salt bridge, which is a concentrated solution

of KCl in agar–agar jelly that completes the electrical circuit and enables the cell to function. Thus, salt bridge provides an electrical contact between the two solutions without allowing them to mix with each other.

The simplest electrochemical cell to study is Daniel cell, which is shown in figure 2. This is prepared by dipping Zn rod in a solution of  $\text{ZnSO}_4$  in one beaker and by dipping Cu strip in a solution of  $\text{CuSO}_4$  in another beaker. It consists of two redox couples,  $\text{Zn}^{2+} | \text{Zn}$  at one end and  $\text{Cu}^{2+} | \text{Cu}$  at the other end. Since Zn has a higher oxidation potential than Cu, so it has a higher tendency to get oxidised than Cu and conversely  $\text{Cu}^{2+}$  has higher tendency to get reduced than  $\text{Zn}^{2+}$ . At this stage, no reaction takes place in either of the two beakers. Now, the two half–cells are connected by connecting wire through an ammeter. As soon as the connection is made, Zn rod starts dissolving i.e. Zn atom changes to  $\text{Zn}^{2+}$  by losing 2 electrons. The  $\text{Zn}^{2+}$  ion passes into the solution, thereby increasing its concentration and 2 electrons remain on the Zn rod. Thus,  $\text{ZnSO}_4$  solution now has +2 unit extra charge and Zn rod has –2 unit extra charge. Thus, there is a charge separation of 4 units between the Zn rod and  $\text{ZnSO}_4$  solution. This charge separation develops a potential referred to as oxidation potential; since oxidation takes place on this electrode. The potential is called standard oxidation potential, if the concentration of  $\text{Zn}^{2+}$  in the solution is 1 M. Similar potential is developed at  $\text{Cu}^{2+} | \text{Cu}$  half–cell also, when  $\text{Cu}^{2+}$  discharges and deposits on Cu rod. This potential is referred to as reduction potential; as reduction takes place on this electrode. The potential is called standard reduction potential, if the concentration of  $\text{Cu}^{2+}$  in the solution is 1 M.






Figure 2. Setup of a Daniel cell

The zinc rod which has the electrons left by the zinc (that got oxidized) becomes negatively charged while the Cu rod which lost electrons to  $\text{Cu}^{2+}$  becomes positively charged. Thus, electric current flows through the connected wire, which is indicated by a deflection in ammeter showing that a chemical reaction is occurring in the cell. During the course of reaction, zinc rod gets dissolved and copper gets deposited on the copper rod. Thus, the concentration of the anode solution increases while that of cathode solution decreases. The flow of electrons occurs from the zinc rod to copper rod in the external circuit and indirectly from cathode to anode in the internal circuit. The current flow is in the direction opposite to electron flow. The reactions occurring at the two electrodes are



This current flow stops after some time. Think, why it happens? As zinc rod dissolves, let one mole of Zn atoms dissolve to form  $\text{Zn}^{2+}$  ions, which pass into the solution and 2 mole of electrons remain on the rod. Thus, rod becomes negatively charged and the solution becomes positively charged. With the passage of time, the solution becomes so much positively charged that any Zn atom getting oxidised and trying to get into the solution, would be repelled by the solution and thus the oxidation of Zn stops. Same phenomenon occurs at cathode and the reduction of  $\text{Cu}^{2+}$  at cathode ceases. As the flow of electrons stop, so does the flow of current. This problem can be removed with the use of salt bridge. The salt bridge contains a highly soluble electrolyte (like KCl,  $\text{NH}_4\text{NO}_3$ ,  $\text{NH}_4\text{Cl}$ ,  $\text{KNO}_3$  etc) in which ionic mobilities of cation and anion are of comparable order.

Now, let us examine the function of salt bridge. Since zinc ions are produced as electrons leave the zinc electrode, this tends to produce a net positive charge in the left compartment. The salt bridge then throws ions having equivalent opposite charge into the solution to maintain electrical neutrality. Thus, the salt bridge keeps the solution neutral by passing appropriate amounts of cations or anions to the two half–cells (compartments).





Thus, the purpose of the salt bridge is to prevent any net charge accumulation in either of the compartments. Salt bridge does not participate chemically in the cell reaction but it is essential for the cell to operate. With the use of salt bridge, the flow of electrons becomes continuous and cell continues to operate. But after some more time, cell ceases to operate. Can you guess why it stops operating now?

## 6.1 cell notation OF AN ELECTROCHEMICAL CELL

- Anode is written on the left side and cathode is written on the right side.
- Phase boundaries are indicated by vertical bar or slash.
- Concentration of the electrolytes in the anode and cathode must be written in parenthesis.
- In case of a gas, the partial pressure is to be mentioned in atm or mm Hg.
- A comma is used to separate two chemical species present in the same solution.
- A double vertical line i.e. || denotes that a salt bridge is present.
- EMF of the cell is written on the extreme right of the representation.

For example,



**Note:** In some cell representations (as in (ii) above), the salt bridge is not indicated which implies that the electrolyte is common to both anode and cathode compartments.

## 7 STANDARD REDUCTION

When the concentrations of the  $\text{Cu}^{2+}$  and  $\text{Zn}^{2+}$  ions are both 1.0 M, we find that the voltage or emf of the Daniell cell is 1.10 V at  $25^\circ\text{C}$ . This voltage must be related directly to the redox reaction, but how? just as the overall cell reaction can be thought of as the sum of two half-cell reactions, the measured emf of the cell can be treated as the sum of the electrical potentials at the Zn and Cu electrodes. Knowing one of these electrode potentials, we could obtain the other by subtraction. It is impossible to measure the potential of just a single electrode, but if we arbitrarily set the potential value of a particular electrode at zero, we can use it to determine the relative potentials of other electrodes. The hydrogen electrode, serves as the reference for this purpose. Hydrogen gas is bubbled into a hydrochloric acid solution at  $25^\circ\text{C}$ . The platinum electrode has two functions. First, it provides a surface on which the dissociation of hydrogen molecules can take place




Second, it serves as an electrical conductor to the external circuit. Under standard state conditions (when the pressure of  $\text{H}_2$  is 1 atm and the concentration of the HCl solution is 1 M), the potential for the reduction of  $\text{H}^+$  at  $25^\circ\text{C}$  is taken to be exactly zero:




The superscript “ $^\circ$ ” denotes standard-state conditions and  $E^\circ$  is **standard reduction potential**, or the voltage associated with a reduction reaction at an electrode when all solutes are 1 M and all gases are at 1 atm. Thus, the standard reduction potential of the hydrogen electrode is defined as zero. The hydrogen electrode is called the standard hydrogen electrode (SHE).

We can use the SHE to measure the potentials of other kinds of electrodes. For example, consider an electrochemical cell with a zinc electrode and a SHE. In this case the zinc electrode is anode and the SHE is the cathode. We deduce this fact from the decrease in mass of the zinc electrode during the operation of the cell, which is consistent with the loss of zinc to the solution caused by the oxidation reaction:




The cell diagram is



As mentioned earlier, the Pt electrode provides the surface on which the reduction takes place. When all the reactants are in their standard states (that is,  $\text{H}_2$  at 1 atm,  $\text{H}^+$  and  $\text{Zn}^{2+}$  ions at



1 M each, the emf of the cell is 0.76 V at 25°C. We can write the half-cell reactions as follows:



By convention, the **standard emf** of the cell,  $E_{\text{cell}}^{\circ}$ , which is composed of a contribution from the anode and a contribution from the cathode, is given by

$$E_{\text{cell}}^{\circ} = E_{\text{cathode}}^{\circ} - E_{\text{anode}}^{\circ}$$


where  $E_{\text{cathode}}^{\circ}$  and  $E_{\text{anode}}^{\circ}$  are the standard reduction potential of the cathode and anode respectively.

For the Zn-SHE cell, we write


$$\begin{aligned} E_{\text{cell}}^{\circ} &= E_{\text{H}^{+}|\text{H}_2}^{\circ} - E_{\text{Zn}^{2+}|\text{Zn}}^{\circ} \\ 0.76 \text{ V} &= 0 - E_{\text{Zn}^{2+}|\text{Zn}}^{\circ} \end{aligned}$$

where the subscript  $\text{H}^{+} | \text{H}_2$  means  $2\text{H}^{+} + 2\text{e}^{-} \longrightarrow \text{H}_2$  and the subscript  $\text{Zn}^{2+} | \text{Zn}$  means  $\text{Zn}^{2+} + 2\text{e}^{-} \longrightarrow \text{Zn}$ . Thus the standard reduction potential of zinc,  $E_{\text{Zn}^{2+}|\text{Zn}}^{\circ}$ , is  $-0.76 \text{ V}$ .


The standard electrode potential of copper can be obtained in a similar fashion, by using a cell with a copper electrode and a SHE. In this case, the copper electrode is the cathode because its mass increases during the operation of the cell, as is consistent with the reduction reaction:



The cell diagram is



and the half-cell reactions are




Under standard-state conditions and at 25°C, the emf of the cell is 0.34 V, so we write

$$\begin{aligned} E_{\text{cell}}^{\circ} &= E_{\text{cathode}}^{\circ} - E_{\text{anode}}^{\circ} \\ 0.34 \text{ V} &= E_{\text{Cu}^{2+}|\text{Cu}}^{\circ} - E_{\text{H}^{+}|\text{H}_2}^{\circ} \\ &= E_{\text{Cu}^{2+}|\text{Cu}}^{\circ} - 0 \end{aligned}$$

In this case the standard reduction potential of copper,  $E_{\text{Cu}^{2+}|\text{Cu}}^{\circ}$ , is 0.34 V, where the subscript  $\text{Cu}^{2+} | \text{Cu}$  means  $\text{Cu}^{2+} + 2\text{e}^{-} \longrightarrow \text{Cu}$ .

Having known the values of standard reduction potential of copper electrode  $E_{\text{Cu}^{2+}|\text{Cu}}^{\circ}$  and that of zinc electrode  $E_{\text{Zn}^{2+}|\text{Zn}}^{\circ}$  it is possible to find out the standard EMF ( $E_{\text{cell}}^{\circ}$ ) of the following electrochemical cell



The standard EMF of the cell is given by

$$\begin{aligned} E_{\text{cell}}^{\circ} &= E_{\text{cathode}}^{\circ} - E_{\text{anode}}^{\circ} \\ &= E_{\text{Cu}^{2+}|\text{Cu}}^{\circ} - E_{\text{Zn}^{2+}|\text{Zn}}^{\circ} \\ &= 0.34 \text{ V} - (-0.76 \text{ V}) \\ &= 1.10 \text{ V}. \end{aligned}$$



## 7.1 ELECTROCHEMICAL SERIES


S. No.	Reduction half cell reaction	E° in volts at 25°C
1.	$F_2 + 2e^- \longrightarrow 2F^-$	+ 2.65
2.	$S_2O_8^{2-} + 2e^- \longrightarrow 2SO_4^{2-}$	+ 2.01
3.	$Co^{3+} + e^- \longrightarrow Co^{2+}$	+ 1.82
4.	$PbO_2 + 4H^+ + SO_4^{2-} + 2e^- \longrightarrow PbSO_4 + 2H_2O$	+ 1.65
5.	$MnO_4^- + 8H^+ + 5e^- \longrightarrow Mn^{2+} + 4H_2O$	+ 1.52
6.	$Au^{3+} + 3e^- \longrightarrow Au$	+ 1.50
7.	$Cl_2 + 2e^- \longrightarrow 2Cl^-$	+ 1.36
8.	$Cr_2O_7^{2-} + 14H^+ + 6e^- \longrightarrow 2Cr^{3+} + 7H_2O$	+ 1.33
9.	$O_2 + 4H^+ + 4e^- \longrightarrow 2H_2O$	+ 1.229
10.	$Br_2 + 2e^- \longrightarrow 2Br^-$	+ 1.07
11.	$NO_3^- + 4H^+ + 3e^- \longrightarrow NO + 2H_2O$	+ 0.96
12.	$2Hg^{2+} + 2e^- \longrightarrow Hg_2^{2+}$	+ 0.92
13.	$Cu^{2+} + I^- + e^- \longrightarrow CuI$	+ 0.86
14.	$Ag^+ + e^- \longrightarrow Ag$	+ 0.799
15.	$Hg_2^{2+} + 2e^- \longrightarrow 2Hg$	+ 0.79
16.	$Fe^{3+} + e^- \longrightarrow Fe^{2+}$	+ 0.77
17.	$I_2 + 2e^- \longrightarrow 2I^-$	+ 0.535
18.	$Cu^+ + e^- \longrightarrow Cu$	+ 0.53
19.	$Cu^{2+} + 2e^- \longrightarrow Cu$	+ 0.34
20.	$Hg_2Cl_2 + 2e^- \longrightarrow 2Hg + 2Cl^-$	+ 0.27
21.	$AgCl + e^- \longrightarrow Ag + Cl^-$	+ 0.222
22.	$Cu^{2+} + e^- \longrightarrow Cu^+$	+ 0.15
23.	$Sn^{4+} + 2e^- \longrightarrow Sn^{2+}$	+ 0.13
24.	$2H^+ + 2e^- \longrightarrow H_2$	0.00
25.	$Fe^{3+} + 3e^- \longrightarrow Fe$	- 0.036
26.	$Pb^{2+} + 2e^- \longrightarrow Pb$	- 0.126
27.	$Sn^{2+} + 2e^- \longrightarrow Sn$	- 0.14
28.	$AgI + e^- \longrightarrow Ag + I^-$	- 0.151
29.	$Ni^{2+} + 2e^- \longrightarrow Ni$	- 0.25
30.	$Co^{2+} + 2e^- \longrightarrow Co$	- 0.28
31.	$Cd^{2+} + 2e^- \longrightarrow Cd$	- 0.403
32.	$Cr^{3+} + e^- \longrightarrow Cr^{2+}$	- 0.41
33.	$Fe^{2+} + 2e^- \longrightarrow Fe$	- 0.44
34.	$Cr^{3+} + 3e^- \longrightarrow Cr$	- 0.74
35.	$Zn^{2+} + 2e^- \longrightarrow Zn$	- 0.762
36.	$2H_2O + 2e^- \longrightarrow H_2 + 2OH^-$	- 0.828
37.	$Mn^{2+} + 2e^- \longrightarrow Mn$	- 1.18
38.	$Al^{3+} + 3e^- \longrightarrow Al$	- 1.66

39.	$\text{H}_2 + 2\text{e}^- \longrightarrow 2\text{H}^-$	- 2.25
40.	$\text{Mg}^{2+} + 2\text{e}^- \longrightarrow \text{Mg}$	- 2.37
41.	$\text{Na}^+ + \text{e}^- \longrightarrow \text{Na}$	- 2.71
42.	$\text{Ca}^{2+} + 2\text{e}^- \longrightarrow \text{Ca}$	- 2.87
43.	$\text{Ba}^{2+} + 2\text{e}^- \longrightarrow \text{Ba}$	- 2.90
44.	$\text{Cs}^+ + \text{e}^- \longrightarrow \text{Cs}$	- 2.92
45.	$\text{K}^+ + \text{e}^- \longrightarrow \text{K}$	- 2.93
46.	$\text{Li}^+ + \text{e}^- \longrightarrow \text{Li}$	- 3.05


The above table lists standard reduction potentials for a number of half-cell reactions.

By definition, the SHE has an  $E^\circ$  value of 0.00 V. Below the SHE the negative standard reduction potentials increase and above it the positive standard reduction potentials increase. It is important to know the following points about the table in calculations:

1. The  $E^\circ$  values apply to the half-cell reactions as read in the forward (left to right) direction.
2. The more positive  $E^\circ$  is, the greater the tendency for the substance to be reduced. For example, the half-cell reaction



has the highest positive  $E^\circ$  value among all the half-cell reactions. Thus,  $\text{F}_2$  is the strongest oxidizing agent because it has the greatest tendency to be reduced. At the other extreme is the reaction



which has the most negative  $E^\circ$  value. Thus  $\text{Li}^+$  is the weakest oxidizing agent because it is the most difficult species to reduce. Conversely, we say that  $\text{F}^-$  is the weakest reducing agent and Li metal is the strongest reducing agent. Under standard-state conditions, the oxidizing agents (the species on the left-hand side of the half-reactions in the given table) increase in strength from bottom to top and the reducing agents (the species on the right-hand side of the half-reactions) increase in strength from top to bottom.


3. The half-cell reactions are reversible. Depending on the conditions, any electrode can act either as an anode or as a cathode. Earlier we saw that the SHE is the cathode ( $\text{H}^+$  is reduced to  $\text{H}_2$ ) when coupled with zinc in a cell and that it becomes the anode ( $\text{H}_2$  is oxidized to  $\text{H}^+$ ) when used in a cell with copper.
4. Under standard-state conditions, any species on the left of a given half-cell reaction will react spontaneously with a species that appears on the right of any half-cell reaction located below it in the given table. This principle is sometimes called the diagonal rule. In the case of the Daniel cell




We see that the substance on the left of the first half-cell reaction is  $\text{Cu}^{2+}$  and the substance on the right in the second half-cell reaction is Zn. Therefore, as we saw earlier, Zn spontaneously reduces  $\text{Cu}^{2+}$  to form  $\text{Zn}^{2+}$  and Cu.

5. Changing the stoichiometric coefficients of a half-cell reaction does not affect the value of  $E^\circ$  because electrode potentials are intensive properties. This means that the value of  $E^\circ$  is unaffected by the size of the electrodes or the amount of solutions present.

For example,




but  $E^\circ$  does not change if we multiply the half-reaction by 2 :



## REDOX REACTIONS AS THE BASIS FOR TITRATION

In acid-base systems we come across with a titration method for finding out the strength of one solution against the other using a pH sensitive indicator. Similarly, in redox systems, the titration method can be adopted to determine the strength of a reductant/ oxidant using a redox sensitive indicator. The usage of indicators in redox titration is illustrated below:

- (i) In one situation, the reagent itself is intensely coloured, e.g. permanganate ion,  $\text{MnO}_4^-$ . Here  $\text{MnO}_4^-$  acts as the self indicator. The visible end point in this case is achieved after the last of the reductant ( $\text{Fe}^{+2}$  or  $\text{C}_2\text{O}_4^{2-}$ ) is oxidised and the first lasting tinge of pink colour appears at  $\text{MnO}_4^-$ , concentration as low as  $10^{-6} \text{ mol dm}^{-3}$ . This ensures a minimal 'overshoot' in colour beyond the equivalence point, the point where the reductant and the oxidant are equal to terms of their mole stoichiometry.
- (ii) If there is no dramatic auto-colour change (as with the  $\text{MnO}_4^-$  titration), there are indicators which are oxidised immediately after the last bit of the reactant is consumed, producing a dramatic colour change.  
Example:  $\text{Cr}_2\text{O}_7^{2-}$  is not a self-indicator, but oxidises the indicator substance diphenylamine just after the equivalence point to produce an intense blue colour, thus signaling the end point.
- (iii) There is yet another method which is interesting and quite common. Its use is restricted to those reagents which are able to oxidise  $\text{I}^-$  ions, say, for example,  $\text{Cu(II)}$ :  $2\text{Cu}^{+2}(\text{aq}) + 4\text{I}^-(\text{aq}) \rightarrow \text{Cu}_2\text{I}_2(\text{s}) + \text{I}_2(\text{aq})$   
This method relies on the facts that iodine itself gives an intense blue colour with starch and has a very specific reaction with thiosulphate ions ( $\text{S}_2\text{O}_3^{2-}$ ), which too is a redox reaction:



$\text{I}_2$ , though insoluble in water, remains in solution containing KI as  $\text{KI}_3$ .

On addition of starch after the liberation of iodine from the reaction of  $\text{Cu}^{+2}$  ions on iodide ions, an intense blue colour appears. This colour disappears as soon as the iodine is consumed by the thiosulphate ions. Thus, the end-point can easily be tracked and the rest is the stoichiometric calculation only.