

Relations & Functions

1 RELATIONS

1.1 INTRODUCTION

In our day to day life, we often talk about relation between two persons, between two straight lines (e.g. perpendicular lines, parallel lines) etc.

Let A be the set of all male students in Delhi whose fathers live in Delhi. Let B be the set of all the people living in Delhi. Let a be a male student living in Delhi i.e. $a \in A$. Let b be the father of a . Then $b \in B$. And a is related to b under son-father relation. If we denote the son-father relation by symbol R then a is related to b under relation R . We can also express this by writing aRb . Here R denotes the relation 'is son of'.

We can also express this statement by saying that the pair of a and b is in relation R i.e., the ordered pair $(a, b) \in R$. This pair (a, b) is ordered in the sense that a and b can't be interchanged because first co-ordinate a represents son, and the second coordinate b represents father of a . Similarly if $a_1 \in A$ and b_1 is father of a_1 , then $(a_1, b_1) \in R$. So we can think of the relation R as a set of ordered pairs whose first coordinate is in A and the second coordinate is in B . Thus $R \subseteq A \times B$. Since the relation 'is son of' i.e., R is a relation relating elements of A to elements of B , we will say that R is a relation from set A to set B .

1.2 DEFINITION

A relation R , from a non-empty set A to another non-empty set B , is a subset of $A \times B$. Equivalently, any subset of $A \times B$ is relation from A to B .

Thus, R is a relation from A to $B \Leftrightarrow R \subseteq A \times B$

$$\Leftrightarrow R \subseteq \{(a, b) : a \in A, b \in B\}$$

Example: Let $A = \{1, 2\}$, $B = \{a, b, c\}$

Let $R = \{(1, a), (1, c)\}$

Here R is a subset of $A \times B$ and hence it is a relation from A to B .

2 DOMAIN AND RANGE OF A RELATION

2.1 DOMAIN OF A RELATION

Let R be a relation from A to B . The domain of relation R is the set of all those elements $a \in A$ such that $(a, b) \in R$ for some $b \in B$. Domain of R is precisely written as domain R .

Thus domain of $(R) = \{a \in A : (a, b) \in R \text{ for some } b \in B\}$

Thus domain of $R =$ set of first components of all the ordered pair which belong to R .

2.2 RANGE OF A RELATION

Let R be a relation from A to B . The range of R is the set of all those elements $b \in B$ such that $(a, b) \in R$ for some $a \in A$.

Thus range of $R = \{b \in B : (a, b) \in R \text{ for some } a \in A\}$.

Range of $R =$ set of second components of all the ordered pairs which belong to R .

Set B is called as codomain of relation R .

Example1: Let $A = \{2, 3, 5\}$ and $B = \{4, 7, 10, 8\}$

Let $aRb \Leftrightarrow a$ divides b

Then $R = (2, 5)$ and range of $R = \{4, 10, 8\}$



Codomain of $R = B = \{4, 7, 10, 8\}$

Example2: Let $A = \{1, 2, 3\}$, $B = \{2, 4, 6, 8\}$

Let R be a relation defined from A to B by $xRy \Leftrightarrow y$ is double of $x, \forall x \in A$

Then $1R2, 2R4, 3R6$

$\therefore R = \{(1, 2), (2, 4), (3, 6)\}$

3 REPRESENTATION OF A RELATION

A relation from a set A to set B can be represented in any one of the following four forms.

3.1 ROSTER FORM

In this form a relation R is represented by the set of all ordered pairs belonging to R .

Example: Let $A = \{-1, 1, 2\}$ and $B = \{1, 4, 9, 10\}$

Let aRb means $a^2 = b$

Then R (in roaster form) = $\{(-1, 1), (1, 1), (2, 4)\}$

3.2 SET-BUILDER FORM

In this form, the relation R is represented as $\{(a, b) : a \in A, b \in B, a \dots b\}$, the blank is to be replaced by the rule which associates a to b .

Example: Let $A = \{1, 3, 5, 7\}$, $B = \{2, 4, 6, 8\}$

Let $R = \{(1, 2), (3, 4), (5, 6), (7, 8)\}$, then R in the builder form can be written as

$R = \{(a, b) : a \in A, b \in B, a - b = -1\}$


3.3 BY ARROW DIAGRAM

In this form, the relation R is represented by drawing arrows from first component to the second component of all ordered pairs belonging to R .

Example: Let $A = \{1, 2, 3, 4\}$, $B = \{0, 2, 4\}$ and R be relation 'is less than' from A to B , then

$R = \{(1, 2), (1, 4), (2, 4), (3, 4)\}$

This relation R from A to B can be represented by the arrow diagram as shown in the figure.



4 TOTAL NUMBER OF RELATIONS

Let A and B be two non empty finite sets having p and q elements respectively.

Then $n(A \times B) = n(A) \cdot n(B) = pq$

Therefore, total number of subsets of $A \times B = 2^{pq}$

Since each subset of $A \times B$ is a relation from A and B , therefore total number of relations form A to B is 2^{pq}

Note: Empty relation ϕ and universal relation $A \times B$ are called trivial relations and any other relation is called a **non trivial relation**.

Example: Let $A = \{1, 2\}$, $B = \{3, 4, 5\}$

Then $n(A \times B) = n(A) \cdot n(B) = 2 \times 3 = 6$

\therefore Number of relations from A to $B = 2^6 = 64$

Important formulae/points

- If R is relation from A to B and $(a, b) \notin R$, then we also write $a \not R b$ (read as a is not related to b)
- In an identity relation on A every element of A should be related to itself only.
- aRb shows that a is the element of domain set and b is the element of range set.

5 FUNCTIONS

The concept of functions is very important because of its close relation with various phenomena of reality. Thus when we square a given real number in fact we perform an operation on the number x to get number x^2 . Hence a function may be viewed as a rule which produces new elements from some given elements. Function is also called mapping or map.

- **Independent Variable**

The symbol which can take an arbitrary value from a given set is called an independent variable.

- **Dependent Variable**

The symbol whose value depends on independent variables is called a dependent variable.

6 DEFINITION OF A FUNCTION

- **Definition 1**

A function f is a relation from a non-empty set A to a non-empty set B such that domain of f is A and no two distinct ordered pairs in f have the same first element.

- **Definition 2**

Let A and B be two non-empty sets, then a rule of which associates each element of A with a unique element of B is called a mapping or a function from A to B we write $f : A \rightarrow B$ (read as f is a function from A to B).

If f associates $x \in A$ to $y \in B$, then we say that y is the image of the element x under the function f or the f image of x and we write $y = f(x)$. The element x is called the pre-image or inverse-image of y .

Thus for a function from A to B :

- A and B should be non-empty.
- Each element of A should have image in B .
- No element of A should have more than one images in B .

7 DOMAIN, CO-DOMAIN AND RANGE OF A FUNCTION

The set A is called as the domain of the map f and the set B is called as the co-domain. The set of the images of all the elements of A under the map f is called the range of f and is denoted by $f(A)$.

Thus range of f i.e. $f(A) = \{f(x) : x \in A\}$.

Clearly $f(A) \subseteq B$

Thus,

- It is necessary that every f image is in B , but there may be some elements in B , which are not f image of any element of A i.e., whose pre-image under f is not in A .
- Two or more elements of A may have same image in B .
- $f : x \rightarrow y$ means that under the function f from A to B , an element x of A has image y in B .
- If domain and range of a function are not to be written, sometimes we denote the function f by writing $y = f(x)$ and read it as y is a function of x .
- A function which has R or one of its subsets as its range is called "real valued function". Further, if its domain is also R or a subset of R , it is called a real function, where R is the set of real numbers.

8 IMPORTANT FUNCTIONS AND THEIR GRAPHS

- **Algebraic functions:** Functions consisting of finite number of terms involving powers and roots of the independent variable with the operations $+$, $-$, \times , \div are called algebraic functions.

Examples: $f(x) = \sqrt{x-1}$, $f(x) = \sqrt{x} + x^3$

- **Polynomial functions:** $f(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$, where $a_0, a_1, a_2, \dots, a_n \in \mathbf{R}$ is said to be a polynomial function of degree n .


- **Logarithmic function:** If $a > 0, a \neq 1$, then the function $y = \log_a x, x \in \mathbf{R}^+$ (set of positive real numbers) is called a logarithmic function, if $a = e$, the logarithmic function is denoted by $\ln x$.

Logarithmic function is the inverse of the exponential function.

For $\log_a x$ to be real, x must be greater than zero.

$$y = \log_a x, a > 0 \text{ and } \neq 1$$


Domain : $(0, \infty)$; Range : $(-\infty, \infty)$;




- **Exponential function:** If $a > 0, a \neq 1$, then the function defined by $y = a^x, x \in \mathbf{R}$ is called an exponential function with base a .

$$y = f(x) = a^x, a > 0, a \neq 1$$

Domain : \mathbf{R} ; Range : $(0, \infty)$;




- **Identity function:** An identity function in x is defined as $f: \mathbf{R} \rightarrow \mathbf{R}, f(x) = x$.



- **Absolute value function:** An absolute value function in x is defined as $f: \mathbf{R} \rightarrow \mathbf{R}, f(x) = |x|$.

$$y = f(x) = |x| = \begin{cases} -x, & x \leq 0 \\ x, & x > 0 \end{cases}$$

Domain : \mathbf{R} ; Range : $[0, \infty)$;



Note that $x = 0$ can be included either with positive values of x or with negative values of x . As we know, all real numbers can be plotted on the real number line, $|x|$ in fact represents the distance of number ' x ' from the origin, measured along the number-line. Thus $|x| \geq 0$. Secondly, any point ' x ' lying on the real number line will have its coordinates as $(x, 0)$. Thus its distance from the origin is $\sqrt{x^2}$. Hence $|x| = \sqrt{x^2}$. Thus we can define $|x|$ as $|x| = \sqrt{x^2}$ e.g. if $x = -2.5$, then $|x| = 2.5$, if $x = 3.8$ then $|x| = 3.8$.

There is another way to define $|x|$ as $|x| = \max \{x, -x\}$.

Basic properties of $|x|$

- $||x|| = |x|$
- Geometrical meaning of $|x - y|$ is the distance between x and y .
- $|x| > a \Rightarrow x > a$ or $x < -a$ if $a \in \mathbf{R}^+$ and $x \in \mathbf{R}$ if $a \in \mathbf{R}^-$.
- $|x| < a \Rightarrow -a < x < a$ if $a \in \mathbf{R}^+$ and $x \in \phi$ if $a \in \mathbf{R}^- \cup \{0\}$
- $|xy| = |x||y|$

- $\left| \frac{x}{y} \right| = \frac{|x|}{|y|}, y \neq 0$

- $|x + y| \leq |x| + |y|$

It is a very useful and interesting property. Here the equality sign holds if x and y either both are non-negative or non-positive (i.e. $x, y \geq 0$). $(|x| + |y|)$ represents the sum of distances of numbers x and y from the origin and $|x + y|$ represents the distance of number $x + y$ from the origin (or distance between ' x ' and ' $-y$ ' measured along the number line).

- $|x - y| \geq |x| - |y|$

Here again the equality sign holds if x and y either both are non-negative or non-positive (i.e. $x, y \geq 0$). $(|x| - |y|)$ represents the difference of distances of numbers x and y from the origin and $|x - y|$ represents the distance between ' x ' and ' y ' measured along the number line.

The last two properties can be put in one compact form i.e., $|x| - |y| \leq |x \pm y| \leq |x| + |y|$.

- **Greatest integer function (step function):** The function $f(x) = [x]$ is called the greatest integer function and is defined as follows:

$[x]$ is the greatest integer less than or equal to x .

Then $[x] = x$ if x is an integer


= integer just less than x if x is not an integer.

Examples: $[3] = 3$, $[2.7] = 2$, $[-7.8] = -8$, $[0.8] = 0$

In other words if we list all the integers less than or equal to x , then the integer greatest among them is called greatest integer of x . Greater integer of x is also called integral part of x .

$$y = f(x) = [x]$$

Domain : \mathbf{R} ; Range : \mathbf{I}




- **Signum function:** The function is defined as

$$y = f(x) = \text{sgn}(x)$$

$$\text{sgn}(x) = \begin{cases} \frac{|x|}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

$$\text{or } \text{sgn}(x) = \begin{cases} -1, & x < 0 \\ 0, & x = 0 \\ 1, & x > 0 \end{cases}$$




Domain : \mathbf{R} ;

Range $\rightarrow \{-1, 0, 1\}$

- **Rational algebraic function:** A function of the form $f(x) = \frac{p(x)}{q(x)}$, where $p(x)$ and $q(x)$ are polynomials and $q(x) \neq 0$, is called a rational function.

The domain of a rational function $\frac{p(x)}{q(x)}$ is the set of all real numbers except points where $q(x) = 0$.

- **Constant function:** The function defined as $f: \mathbf{R} \rightarrow \{c\}$ where $f(x) = c$



9 ALGEBRAIC OPERATIONS ON FUNCTIONS

Let us consider two functions.

$f: D_1 \rightarrow \mathbf{R}$ and $g: D_2 \rightarrow \mathbf{R}$. We describe functions $f + g$, $f - g$, $f \cdot g$ and f/g as follows:

- $f + g: D \rightarrow \mathbf{R}$ is a function defined by $(f + g)x = f(x) + g(x)$, where $D = D_1 \cap D_2$
- $f - g: D \rightarrow \mathbf{R}$ is a function defined by $(f - g)x = f(x) - g(x)$, where $D = D_1 \cap D_2$
- $f \cdot g: D \rightarrow \mathbf{R}$ is a function defined by $(f \cdot g)x = f(x) \cdot g(x)$, where $D = D_1 \cap D_2$
- $f/g: D \rightarrow \mathbf{R}$ is a function defined by $(f/g)x = \frac{f(x)}{g(x)}$, where $D = D_1 \cap \{x \in D_2 : g(x) \neq 0\}$
- $(\alpha f)(x) = \alpha f(x)$, $x \in D_1$ and α is any real number.

10 TYPES OF FUNCTIONS

We have seen that f is a function from A to B , if each element of A has image in B and no element of A has more than one images in B .

But for a function f from A to B following possibilities are there


- Distinct elements of A have distinct images in B .
- More than one element of A may have same image in B .
- Each element of B is the image of some element of A .
- There may be some elements in B which are not the images of any element of A .

Because of the above mentioned possibilities, we have the following types of functions:

10.1 One-one or injective map

A map $f: A \rightarrow B$ is said to be one-one or injective if each and every element of set A has distinct images in set B .

The map $f: A\{-1, 1, 3\} \rightarrow B\{-2, 2, 6, 7\}$ given by $f(x) = 2x$ is a one-one map.






10.2 Many one map:


A map $f : A \rightarrow B$ is said to be many one if and only if it is not one-one.

The map $f : A \rightarrow B$ given by $f(x) = x^2$ is a many-one map.



10.3 Onto map or surjective map:

A map $f : A \rightarrow B$ is said to be onto map or surjective map if and only if each element of B is the image of some element of A i.e. if and only if for every $y \in B$ there exists some $x \in A$ such that $y = f(x)$.



Thus f is onto iff $f(A) = B$ i.e. range of $f =$ co-domain of f .


A map $f : A \rightarrow B$ given by $f(x) = 3x$ is an onto map.

Note: Functions which are not onto, are into.

10.4 One-one onto map or bijective map:

A map $f : A \rightarrow B$ is said to be one-one onto or bijective if and only if it is both one-one and onto i.e., if

- distinct element of A have distinct images in B .
- each element of B is the image of some element of A .



The map $f : A \rightarrow B$ given by $f(x) = 2x$ is a one-one onto map.

- A one-one onto function is also called a one-to-one correspondence or one-one correspondence.
- Let $f : A \rightarrow B$ be a function from finite set A to finite set B . Then
 - f is one-one $\Rightarrow n(A) \leq n(B)$
 - f is onto $\Rightarrow n(B) \leq n(A)$
 - f is one-one onto $\Rightarrow n(A) = n(B)$

11 COMPOSITION OF FUNCTIONS


Let A, B, C be three non-empty sets, f be a function from A to B and g be a function from B to C . The question arises : can we combine these two functions to get a new function? Yes! The most natural way of doing this is to send every element $x \in A$ in two stages to an element of C ; first by applying f to x and then by applying g to the resulting element $f(x)$ of B .

DEFINITION

Let $f : A \rightarrow B$ and $g : B \rightarrow C$ be any two mappings. Then f maps an element $x \in A$ to an element $f(x) = y \in B$ and this y is mapped by g to an element $z \in C$. Thus $z = g(y) = g(f(x))$

Thus we have a rule, which associates with each $x \in A$, a unique element $z = g(f(x))$ of C . This rule is therefore a mapping from A to C . We denote this mapping by $g \circ f$ (read as 'g composition f') and call it the composite mapping of f and g .

$$(g \circ f)(x) = g(f(x)) \quad \forall x \in A$$



The composition of two functions is also called the resultant of two functions or the function of a function.

Observe that the order of events occur from right to left i.e. gof reads composite of f and g and it means that we have to first apply f and then follow it up with g .

Note that for the composite function gof to exist, it is essential that range of f must be a subset of domain of g .

- (i) $\text{Dom. } (gof) = \{x : x \in \text{domain } (f), f(x) \in \text{domain } (g)\}$
- (ii) If gof is defined then it is not necessary that fog is defined.

12 INVERSE FUNCTION

Let f be one-one and onto map from A to B . Since f is onto, therefore $\forall y \in B$ there exist $x \in A$ such that $f(x) = y$ and since f is one-one therefore this element x is unique. Thus we can define a map, say g from B onto A such that $g(y) = x$. This map g is called inverse map of f and is denoted by f^{-1} .

Thus $f^{-1} : B \rightarrow A$ such that $f^{-1}(y) = x$ iff $f(x) = y$


- **How to find the inverse of a given function?**

In order to find the inverse of the function $f(x)$, let $y = f(x)$

From this express x in terms of y . This value of x in terms of y will be $f^{-1}(y)$. Now put x in place of y in $f^{-1}(y)$ to get $f^{-1}(x)$.

Note: f^{-1} exists if and only if f is one-one onto.

$$\text{Let } y = f(x) \Rightarrow y = 2x \Rightarrow x = \frac{y}{2} \Rightarrow f^{-1}(y) = \frac{y}{2} \Rightarrow f^{-1}(x) = \frac{x}{2}$$



Important formulae/points

Function f from A to B have the properties.

- Distinct elements of A may have distinct images in B .
- More than one element of A may have same image in B .
- There may be some elements in B which are not the images of any element of A .